



DIFFEOLOGICAL MORITA EQUIVALENCE

Nesta VAN DER SCHAAF

Résumé. Nous introduisons une nouvelle notion d'équivalence de Morita pour les groupoïdes difféologiques, généralisant la notion originale pour les groupoïdes de Lie. Pour cela, nous développons une théorie des actions de groupoïdes difféologiques, fibrés et bi-fibrés. Nous définissons une notion de fibré principal qui utilise la notion de subduction, généralisant la notion de fibré principal pour un group(oid) de Lie. Nous disons que deux groupoïdes difféologiques sont Morita équivalents si, et seulement si, il existe un fibré bi-principal entre eux. Utilisant le produit tensoriel de Hilsum-Skandalis, nous définissons en outre une composition des bi-fibrés difféologiques, et obtenons une bi-catégorie **DiffeolBiBund**. Notre principal résultat est le suivant: un bi-fibré est bi-principal si, et seulement si, il est faiblement inversible dans cette bi-catégorie. Ceci généralise un théorème bien connu de la théorie des groupoïdes de Lie. Comme application, nous prouvons que les espaces d'orbites de deux groupoïdes difféologiques Morita équivalents sont difféomorphes. Nous montrons également que les propriétés d'un groupoïde difféologique d'être fibrant, et sa catégorie d'actions, sont des invariants de Morita.

Abstract. We introduce a new notion of *Morita equivalence* for *diffeological groupoids*, generalising the original notion for Lie groupoids. For this we develop a theory of *diffeological groupoid actions*, *-bundles* and *-bibundles*. We define a notion of *principality* for these bundles, which uses the notion of a *subduction*, generalising the notion of a Lie group(oid) principal bundle.

We say two diffeological groupoids are Morita equivalent if and only if there exists a *biprincipal* bibundle between them. Using a Hilsum-Skandalis tensor product, we further define a composition of diffeological bibundles, and obtain a bicategory **DiffeolBiBund**. Our main result is the following: a bibundle is biprincipal if and only if it is *weakly invertible* in this bicategory. This generalises a well known theorem from the Lie groupoid theory. As an application of the framework, we prove that the *orbit spaces* of two Morita equivalent diffeological groupoids are diffeomorphic. We also show that the property of a diffeological groupoid to be *fibrating*, and its *category of actions*, are Morita invariants.

Keywords. Diffeology, Lie groupoids, diffeological groupoids, bibundles, Hilsum-Skandalis products, Morita equivalence, orbit spaces.

Mathematics Subject Classification (2010). 22Axx, 22A22, 58H05.

1. Introduction

Diffeology originates from the work of J.-M. Souriau [Sou80; Sou84] and his students [DI83; Don84; Igl85] in the 1980s. The main objects of this theory are *diffeological spaces*, a type of generalised smooth space that extends the traditional notion of a smooth manifold. They make for a convenient framework that deals well with (singular) quotients, function spaces (or otherwise infinite-dimensional objects), fibred products (or otherwise singular subspaces), and other constructions that lie beyond the realm of classical differential topology. As many of these constructions naturally occur in differential topology and -geometry, and since they cannot be studied with their standard tools, diffeology has become a useful addition to the geometer's toolbox.

Diffeological groupoids have recently garnered attention in the mathematical physics of general relativity [BFW13; Gl19], foliation theory [ASZ19; GZ19; Mac20], the theory of algebroids [AZ], the theory of (differentiable) stacks [RV18; WW19], and even in relation to noncommutative geometry [IZL18; IZP20]. In all but one of these fields (general relativity), the notion of *Morita equivalence* is an important one. Yet, as the authors of [GZ19, p.3] point out: “The theory of Morita equivalence for diffeological groupoids has not been developed yet.” In the current paper we present one possible development of such a notion, based on the results of the author's Mas-

ter thesis [vdS20]. This development is a generalisation of the theory of Hilsum-Skandalis bibundles and the Morita equivalence of Lie groupoids, where many definitions and proofs, and certainly the general idea, extend quite straightforwardly to the diffeological case. The main exception is that we need to replace surjective submersions with so-called *subductions*. This special type of smooth map is, even on smooth manifolds, slightly weaker than the notion of a surjective submersion, but it turns out that they still share enough of their properties so that the entire theory can be developed¹. This development proceeds roughly as follows: based on the notions of *actions* and *bundles* defined in Section 4, we define a diffeological version of a *bibundle* between groupoids (Definition 5.1). These stand in analogy to *bimodules* for rings, and can be treated as a generalised type of morphism between groupoids. This gives a *bicategory* $\mathbf{DiffeolBiBund}$ of diffeological groupoids, bibundles, and *biequivariant maps* (Theorem 5.17). Using the aforementioned notion of a *subduction* (Definition 2.16), we define *biprincipality* of bibundles, and with this, we obtain a notion of *Morita equivalence* for diffeological groupoids (Definition 5.3). In the bicategory we also get a notion of equivalence, by way of the *weak isomorphisms*. A morphism in a bicategory is called *weakly invertible* if it is invertible *up to 2-isomorphism*. Two objects in a bicategory are called *weakly isomorphic* if there exists a weakly invertible morphism between them. The main point of this paper is to prove a *Morita theorem* for diffeological groupoids, characterising the weakly invertible bibundles, and hence realising Morita equivalence as a particular instance of weak isomorphism:

Theorem 5.31 (Morita theorem). *A diffeological bibundle is weakly invertible if and only if it is biprincipal. In other words, two diffeological groupoids are Morita equivalent if and only if they are weakly isomorphic in the bicategory $\mathbf{DiffeolBiBund}$.*

A Morita theorem for Lie groupoids has been known in the literature for some time, see e.g. [Lan01b, Proposition 4.21]. Throughout the paper, we shall point out some differences between the diffeological- and Lie theories. The main difference is that, due to technical constraints, a Morita theorem for Lie groupoids only holds in the restricted setting of *left principal* bibundles.

¹This is essentially due to the fact that the subductions are the *strong epimorphisms* in the category of diffeological spaces [BH11, Proposition 5.10].

The main improvement of Theorem 5.31 over the classical Lie Morita theorem, besides the generalisation to diffeology, is therefore that it considers also a more general class of bibundles. Besides this improvement, with this paper we hope to contribute a complete account of the basic theory of bibundles and Morita equivalence of groupoids, providing detailed proofs and constructions of most necessary technical results, and culminating in a proof of the main Theorem 5.31. A brief outline of the contents of the paper is as follows.

We briefly recall the definition of a diffeology in Section 2. In particular, we describe the diffeologies of fibred products (pullbacks) and quotients, since they will be important to describe the smooth structure of the orbit space and space of composable arrows of a groupoid. We also define and study the behaviour of *subductions*, especially in relation to fibred products.

In Section 3 we define *diffeological groupoids*, and highlight some examples from the literature.

Sections 4 and 5 contain the main contents of this paper. In them, we define the notions of smooth groupoid *actions* and *-bundles*. For the latter we give a new notion of *principality*, generalising the notion of a principal Lie group(oid) bundle. This leads naturally to the definition of a *biprincipal bibundle*, and hence to our definition of *Morita equivalence*. The remainder of Section 5 is dedicated to a proof of Theorem 5.31.

In Section 6, we describe some *Morita invariants*, by generalising some well-known theorems from the Lie theory. We prove: the property of a diffeological groupoid to be *fibrating* is preserved under our notion of Morita equivalence; the *orbit spaces* of two Morita equivalent diffeological groupoids are diffeomorphic; and the categories of representations of two Morita equivalent diffeological groupoids are categorically equivalent.

Lastly, in Section 7, we discuss the question of diffeological Morita equivalence between Lie groupoids. We end the paper with the open Question 7.6, and some suggestions for future research.

Acknowledgements. The author thanks Klaas Landsman and Ioan Mărcuț for being the supervisor and second reader of his Master thesis, respectively, and for encouraging him to write the current paper! He also thanks Klaas for feedback on an earlier version of the paper, and Patrick Iglesias-Zemmour for email correspondence. Lastly, he thanks the anonymous referee for their useful and thoughtful feedback, which helped to improve the clarity of the exposition.

2. Diffeology

One of the main conveniences of *diffeology*² is that the category **Diffeol** of diffeological spaces and smooth maps (Definition 2.2) is complete, cocomplete, (locally) Cartesian closed, and in fact a quasitopos [BH11, Theorem 3.2]. This means that we can perform many categorical constructions that are unavailable in the category **Mnfd** of smooth manifolds. From these, the ones that are important for us are pullbacks and quotients. We discuss both of these explicitly below. The approach of diffeology has been compared to other theories of generalised smooth spaces in [Sta11; BIKW17]. For some historical remarks we refer to [IZ13b; IZ17] and [vdS20, Chapter I]. The main reference for this section is the textbook [IZ13a] by Iglesias-Zemmour, in which nearly all of the theory below is already developed.

Definition 2.1. A *parametrisation on a set X* is a function $U \rightarrow X$ defined on an open subset $U \subseteq \mathbb{R}^m$ of Euclidean space, for arbitrary $m \in \mathbb{N}_{\geq 0}$. We denote by $\text{Param}(X)$ the set of all parametrisations on X .

The basic idea behind diffeology is that it determines which parametrisations are ‘smooth’, in such a way that it captures the properties of ordinary smooth functions on smooth manifolds. The precise definition is as follows:

Definition 2.2 (Axioms of Diffeology). *Let X be a set. A diffeology on X is a collection of parametrisations $\mathcal{D}_X \subseteq \text{Param}(X)$, containing what we call plots, satisfying the following three axioms:*

- (Covering) *Every constant parametrisation $U \rightarrow X$ is a plot.*
- (Smooth Compatibility) *For every plot $\alpha : U_\alpha \rightarrow X$ in $\mathcal{D}_X$ and every smooth function $h : V \rightarrow U_\alpha$ between open subsets of Euclidean space, we have that $\alpha \circ h \in \mathcal{D}_X$.*
- (Locality) *If $\alpha : U_\alpha \rightarrow X$ is a parametrisation, and $(U_i)_{i \in I}$ an open cover of U_α such that each restriction $\alpha|_{U_i}$ is a plot of X , then $\alpha \in \mathcal{D}_X$.*

²The etymology of the word is explained in the afterword to [IZ13a]. Souriau first used the term “*différentiel*”, as in ‘differential’ (from the Latin *differentia*, “difference”). Through a suggestion by Van Est, the name was later changed to “*difféologie*,” as in “*topologie*” (‘topology’, from the Ancient Greek *tópos*, “place,” and *-(o)logy*, “study of”). Hence the term: diffeology.

A set X , paired with a diffeology: $(X, \mathcal{D}_X)$, is called a diffeological space. Although, usually we shall just write X .

A function $f : (X, \mathcal{D}_X) \rightarrow (Y, \mathcal{D}_Y)$ between diffeological spaces is called smooth if for every plot $\alpha \in \mathcal{D}_X$ of X , the composition $f \circ \alpha \in \mathcal{D}_Y$ is a plot of Y . The set of all smooth functions between such diffeological spaces is denoted $C^\infty(X, Y)$, and smoothness is preserved by composition. The category of diffeological spaces and smooth maps is denoted by **Diffeol**, and the isomorphisms in this category are called diffeomorphisms.

Example 2.3. Any open subset $U \subseteq \mathbb{R}^m$ of Euclidean space, for $m \in \mathbb{N}_{\geq 0}$, gets a canonical diffeology $\mathcal{D}_U$, called the *Euclidean diffeology*. Its plots are the parametrisations that are smooth in the ordinary sense of the word. Similarly, we get a canonical diffeology $\mathcal{D}_M$ for any smooth manifold M , called the *manifold diffeology*. With respect to these diffeologies, the notion of smoothness defined in Definition 2.2 agrees with the ordinary one. Hence the inclusion functor $\mathbf{Mnfd} \hookrightarrow \mathbf{Diffeol}$ is fully faithful, and we can adopt the previous definition without causing any confusion.

Example 2.4. Any set X carries two canonical diffeologies. First, the largest diffeology, $\mathcal{D}_X^\bullet := \text{Param}(X)$, called the *coarse diffeology*, containing all possible parametrisations. Letting $X^\bullet$ denote the diffeological space with the coarse diffeology, it is easy to see that every function $Z \rightarrow X^\bullet$ is smooth. On the other hand, the smallest diffeology on X is $\mathcal{D}_X^\circ$, containing all locally constant parametrisations. This is called the *discrete diffeology*. Similar to the above, we find that every function $X^\circ \rightarrow Y$ is smooth.

Example 2.5. For any two diffeological spaces X and Y , there is a natural diffeology on the space of smooth functions $C^\infty(X, Y)$ called the *standard functional diffeology* [IZ13a, Article 1.57]. It is the smallest diffeology that makes the evaluation map $(f, x) \mapsto f(x)$ smooth. With these diffeologies, **Diffeol** becomes Cartesian closed.

2.1 Generating families

The Axiom of Locality in Definition 2.2 ensures that the smoothness of a parametrisation, or of a function between diffeological spaces, can be checked locally. This allows us to introduce the following notions, which will help us study interesting constructions, and will often simplify proofs.

Definition 2.6. Consider a family $\mathcal{F} \subseteq \text{Param}(X)$ of parametrisations on X . There exists a smallest diffeology on X that contains $\mathcal{F}$. We denote this diffeology by $\langle \mathcal{F} \rangle$, and call it the diffeology generated by $\mathcal{F}$. If $\mathcal{D}_X = \langle \mathcal{F} \rangle$, we say $\mathcal{F}$ is a generating family for $\mathcal{D}_X$. The elements of $\mathcal{F}$ are called generating plots.

The plots of the diffeology generated by $\mathcal{F}$ are characterised as follows: a parametrisation $\alpha : U_\alpha \rightarrow X$ is a plot in $\langle \mathcal{F} \rangle$ if and only if α is locally either constant, or factors through elements of $\mathcal{F}$. Concretely, this means that for all $t \in U_\alpha$ there exists an open neighbourhood $t \in V \subseteq U_\alpha$ such that $\alpha|_V$ is either constant, or of the form $\alpha|_V = F \circ h$, where $F : W \rightarrow X$ is an element in $\mathcal{F}$, and $h : V \rightarrow W$ is a smooth function between open subsets of Euclidean space. When the family $\mathcal{F}$ is *covering*, in the sense that $\bigcup_{F \in \mathcal{F}} \text{im}(F) = X$, then the condition for $\alpha|_V$ to be constant becomes redundant, and the plots in $\langle \mathcal{F} \rangle$ are locally just of the form $\alpha|_V = F \circ h$.

The main use of this construction is that we may encounter families of parametrisations that are not quite diffeologies, but that contain functions that we nevertheless want to be smooth. On the other hand, calculations may sometimes be simplified by finding a suitable generating family for a given diffeology. This simplification lies in the following result, saying that smoothness has only to be checked on generating plots:

Proposition 2.7. Let $f : X \rightarrow Y$ be a function between diffeological spaces, such that $\mathcal{D}_X$ is generated by some family $\mathcal{F}$. Then f is smooth if and only if for all $F \in \mathcal{F}$ we have $f \circ F \in \mathcal{D}_Y$.

Example 2.8. The *wire diffeology* (called the *spaghetti diffeology* by Souriau) is the diffeology $\mathcal{D}_{\text{wire}}$ on $\mathbb{R}^2$ generated by $C^\infty(\mathbb{R}, \mathbb{R}^2)$. The resulting diffeological space is not diffeomorphic to the ordinary $\mathbb{R}^2$, since the identity map $\text{id}_{\mathbb{R}^2} : (\mathbb{R}^2, \mathcal{D}_{\mathbb{R}^2}) \rightarrow (\mathbb{R}^2, \mathcal{D}_{\text{wire}})$ is not smooth.

Example 2.9. The charts of a smooth atlas on a manifold define a generating family for the manifold diffeology from Example 2.3. Since a manifold may have many atlases, this shows that similarly any diffeology may have many generating families.

2.2 Quotients

We use the terminology from Section 2.1 to define a natural diffeology on a quotient $X/\sim$. This question relates to a more general one: given a function $f : X \rightarrow Y$, and a diffeology $\mathcal{D}_X$ on the domain, what is the smallest diffeology on Y such that f remains smooth? The following provides an answer:

Definition 2.10. *Let $f : X \rightarrow Y$ be a function between sets, and let $\mathcal{D}_X$ be a diffeology on X . The pushforward diffeology on Y is the diffeology $f_*(\mathcal{D}_X) := \langle f \circ \mathcal{D}_X \rangle$, where $f \circ \mathcal{D}_X$ is the family of parametrisations of the form $f \circ \alpha$, for $\alpha \in \mathcal{D}_X$. The pushforward diffeology is the smallest diffeology on Y that makes f smooth.*

We can now use this to define a natural diffeology on a quotient space:

Definition 2.11. *Let X be a diffeological space, and let $\sim$ be an equivalence relation on the set X . We denote the equivalence classes by $[x]$. The quotient $X/\sim$ is the collection of all equivalence classes, and comes with a canonical projection map $p : X \rightarrow X/\sim$, which sends $x \mapsto [x]$. The quotient diffeology on $X/\sim$ is defined as the pushforward diffeology $p_*(\mathcal{D}_X)$ of $\mathcal{D}_X$ along the canonical projection map. Naturally, with respect to this diffeology, the canonical projection map becomes smooth.*

The quotient diffeology will be used extensively, where the equivalence relation will often be defined by the orbits of a group(oid) action, or as the fibres of some smooth surjection. The existence of the quotient diffeology for arbitrary quotients should be contrasted to the situation for smooth manifolds, where quotients often carry no natural differentiable structure at all, but where instead one could appeal to the *Godement criterion* ([Ser65, Theorem 2, p. 92]). The following is an example of a quotient that does not exist as a smooth manifold, but whose diffeological structure is still quite rich:

Example 2.12. The *irrational torus* is the diffeological space defined by the quotient of $\mathbb{R}$ by an additive subgroup: $T_\theta := \mathbb{R}/(\mathbb{Z} + \theta\mathbb{Z})$, where $\theta \in \mathbb{R} \setminus \mathbb{Q}$ is an arbitrary irrational number. Equivalently, it can be described as the leaf space of the Kronecker foliation on the 2-torus with irrational slope. The topology of this quotient contains only the two trivial open sets, yet its quotient

diffeology is non-trivial³. They were first classified in [DI83], whose result is (amazingly) directly analogous to the classification of the irrational rotation algebras [Rie81]. This example is treated in detail in [vdS20, Section 2.3].

2.3 Fibred products

The second construction we need is that of *fibred products*, which are the pullbacks in the category **Diffeol**. Recall that if $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ are two functions between sets with a common codomain, then the fibred product of sets is (up to unique bijection)

$$X \times_Z^{f,g} Y := \{(x, y) \in X \times Y : f(x) = g(y)\}.$$

When each set is equipped with a diffeology, we shall construct a diffeology on the fibred product in two steps. First we describe a natural diffeology on the product $X \times Y$, and then show how this descends to a diffeology on the fibred product as a subset.

Definition 2.13. *Let X and Y be two diffeological spaces. The product diffeology on the Cartesian product $X \times Y$ is defined as*

$$\mathcal{D}_{X \times Y} := \langle \mathcal{D}_X \times \mathcal{D}_Y \rangle,$$

where $\mathcal{D}_X \times \mathcal{D}_Y$ is the family of parametrisations of the form $\alpha_1 \times \alpha_2$, for $\alpha_1 \in \mathcal{D}_X$ and $\alpha_2 \in \mathcal{D}_Y$. The plots in $\mathcal{D}_{X \times Y}$ are exactly the parametrisations $\alpha : U_\alpha \rightarrow X \times Y$ such that $\text{pr}_1 \circ \alpha$ and $\text{pr}_2 \circ \alpha$ are plots of X and Y , respectively. We assume that products are always furnished with their product diffeologies.

It is clear that both projection maps pr_1 and pr_2 are smooth with respect to the product diffeology. The smooth functions into $X \times Y$ behave exactly as one would expect, where $f : A \rightarrow X \times Y$ is smooth if and only if the components $f_1 = \text{pr}_1 \circ f$ and $f_2 = \text{pr}_2 \circ f$ are smooth.

Next we define how the diffeology on a set X transfers to any of its subsets:

³This shows that there are meaningful notions of smooth space that do not rely on the regnant philosophy of “smooth space = topological space + extra structure.”

Definition 2.14. Consider a diffeological space X , and an arbitrary subset $A \subseteq X$. Let $i_A : A \hookrightarrow X$ denote the natural inclusion map. The subset diffeology on A is defined as

$$\mathcal{D}_{A \subseteq X} := \{\alpha \in \text{Param}(A) : i_A \circ \alpha \in \mathcal{D}_X\}.$$

That is, α is a plot of A if and only if, when seen as a parametrisation of X , it is also a plot. We assume that a subset of a diffeological space is always endowed with its subset diffeology.

Since the fibred product $X \times_Z^{f,g} Y$ is a subset of the product $X \times Y$, the following definition is a natural combination of Definitions 2.13 and 2.14:

Definition 2.15. Let $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ be two smooth maps between diffeological spaces. The fibred product diffeology $\mathcal{D}_{X \times_Z^{f,g} Y}$ on the set $X \times_Z^{f,g} Y$ is the subset diffeology it gets from the product diffeology on $X \times Y$. Concretely:

$$\mathcal{D}_{X \times_Z^{f,g} Y} = \{\alpha \in \mathcal{D}_{X \times Y} : f \circ \alpha_1 = g \circ \alpha_2\}.$$

That is, the plots of the fibred product are just plots of $X \times Y$, whose components satisfy an extra condition. We assume that all fibred products are equipped with their fibred product diffeologies.

2.4 Subductions

Subductions are a special class of smooth functions that generalise the notion of surjective submersion from the theory of smooth manifolds. Since there is no unambiguous notion of tangent space in diffeology (cf. [CW16]), the definition looks somewhat different. For (more) detailed proofs of the results in this section, we refer to [IZ13a, Article 1.46] and surrounding text, and [vdS20, Section 2.6].

Definition 2.16. A surjective function $f : X \rightarrow Y$ between diffeological spaces is called a subduction if $f_*(\mathcal{D}_X) = \mathcal{D}_Y$. Note that subductions are automatically smooth.

In the case that f is a subduction, since it is then particularly a surjection, the family of parametrisations $f \circ \mathcal{D}_X$ is covering, and hence the plots of $\mathcal{D}_Y$

are all locally of the form $f \circ \alpha$, where $\alpha \in \mathcal{D}_X$. In other words, f is a subduction if and only if f is smooth and the plots of Y can locally be *lifted* along f to plots of X :

Lemma 2.17. *Let $f : X \rightarrow Y$ be a function between diffeological spaces. Then f is a subduction if and only if the following two conditions are satisfied:*

1. *The function f is smooth.*
2. *For every plot $\alpha : U_\alpha \rightarrow Y$, and any point $t \in U_\alpha$, there exists an open neighbourhood $t \in V \subseteq U_\alpha$ and a plot $\beta : V \rightarrow X$, such that $\alpha|_V = f \circ \beta$.*

Since many of the functions we encounter will naturally be smooth already, the notion of subductiveness is effectively captured by condition (2) in this lemma. This can also be seen in the following simple example:

Example 2.18. Consider the product $X \times Y$ of two diffeological spaces X and Y . The projection maps pr_1 and pr_2 are both subductions.

Example 2.19. For a surjective function $\pi : X \rightarrow B$ we get an equivalence relation on X , where two points are identified if and only if they inhabit the same π -fibre. The equivalence classes are exactly the π -fibres themselves. We denote the quotient set of this equivalence relation by X/π , and equip it with the quotient diffeology whenever X is a diffeological space. If π is a subduction, then there is a diffeomorphism $B \cong X/\pi$ [IZ13a, Article 1.52].

For subsequent use, we state here some useful properties of subductions with respect to composition:

Lemma 2.20. *We have the following properties for subductions:*

1. *If f and g are two subductions, then the composition $f \circ g$ is a subduction as well.*
2. *Let $f : Y \rightarrow Z$ and $g : X \rightarrow Y$ be two smooth maps such that the composition $f \circ g$ is a subduction. Then so is f .*
3. *Let $\pi : X \rightarrow B$ be a subduction, and $f : B \rightarrow Y$ an arbitrary function. Then f is smooth if and only if $f \circ \pi$ is smooth. In fact, f is a subduction if and only if $f \circ \pi$ is a subduction.*

Proof. (1) This is [IZ13a, Article 1.47].

(2) Assume $f : Y \rightarrow Z$ and $g : X \rightarrow Y$ are smooth, such that $f \circ g$ is a subduction. Take a plot $\alpha : U_\alpha \rightarrow Z$. Since the composition is a subduction, for every $t \in U_\alpha$ we can find an open neighbourhood $t \in V \subseteq U_\alpha$ and a plot $\beta : V \rightarrow X$ such that $\alpha|_V = (f \circ g) \circ \beta$. Since g is smooth, we get a plot $g \circ \beta \in \mathcal{D}_Y$, which is a local lift of α along f . The result follows by Lemma 2.17.

(3) If f is smooth, it follows immediately that $f \circ \pi$ is smooth. Suppose now that $f \circ \pi$ is smooth. We need to show that f is smooth. For that, take a plot $\alpha : U_\alpha \rightarrow B$. Since π is a subduction, we can find an open cover $(V_t)_{t \in U_\alpha}$ of U_α together with a family of plots $\beta_t : V_t \rightarrow X$ such that $\alpha|_{V_t} = \pi \circ \beta_t$. It follows that each restriction $f \circ \alpha|_{V_t} = f \circ \pi \circ \beta_t$ is smooth, and by the Axiom of Locality it follows that $f \circ \alpha \in \mathcal{D}_Y$, and hence that f is smooth. The claim about when f is a subduction follows from (2). $\square$

We also collect the following noteworthy claim:

Proposition 2.21 ([IZ13a, Article 1.49]). *An injective subduction is a diffeomorphism.*

We recall now some elementary results on the interaction between subductions and fibred products, as obtained in [vdS20, Section 2.6]. We point out that if f is a subduction, an arbitrary restriction $f|_A$ may no longer be a subduction. We know from Example 2.18 that the second projection map pr_2 of a product $X \times Y$ is a subduction, but it is not always the case that the restriction of this projection to a fibred product $X \times_Z^{f,g} Y$ is a subduction as well. The following result shows that, to ensure this, it suffices to assume that f is a subduction:

Lemma 2.22. *Let $f : X \rightarrow Z$ be a subduction, and let $g : Y \rightarrow Z$ be a smooth map. Then the restricted projection map*

$$\text{pr}_2|_{X \times_Z^{f,g} Y} : X \times_Z^{f,g} Y \longrightarrow Y$$

*is also a subduction. In other words, in **Diffeol**, subductions are preserved under pullback.*

Proof. Consider a plot $\alpha : U_\alpha \rightarrow Y$. By composition, this gives another plot $g \circ \alpha \in \mathcal{D}_Z$. Now, since f is a subduction, for every $t \in U_\alpha$ we can find a

plot $\beta : V \rightarrow X$ defined on an open neighbourhood $t \in V \subseteq U_\alpha$ such that $g \circ \alpha|_V = f \circ \beta$. This gives a plot $(\beta, \alpha|_V) : V \rightarrow X \times_Z Y$ that satisfies $\text{pr}_2|_{X \times_Z Y} \circ (\beta, \alpha|_V) = \alpha|_V$. The result follows by Lemma 2.17. $\square$

The next result shows how two subductions interact with fibred products:

Lemma 2.23. *Consider the following two commuting triangles of diffeological spaces and smooth maps:*

$$\begin{array}{ccc} X_1 & \xrightarrow{f} & Y_1 \\ & \searrow r & \swarrow R \\ & A & \end{array} \quad \text{and} \quad \begin{array}{ccc} X_2 & \xrightarrow{g} & Y_2 \\ & \searrow l & \swarrow L \\ & A, & \end{array}$$

where both f and g are subductions. Then the map

$$(f \times g)|_{X_1 \times_A X_2} : X_1 \times_A^{r,l} X_2 \longrightarrow Y_1 \times_A^{R,L} Y_2; \quad (x_1, x_2) \longmapsto (f(x_1), g(x_2))$$

is also a subduction.

Proof. Clearly $f \times g$ is smooth, so we are left to show that the second condition in Lemma 2.17 is fulfilled. For that, take a plot

$$(\alpha_1, \alpha_2) : U \longrightarrow Y_1 \times_A^{R,L} Y_2,$$

i.e., we have two plots $\alpha_1 \in \mathcal{D}_{Y_1}$ and $\alpha_2 \in \mathcal{D}_{Y_2}$ such that $R \circ \alpha_1 = L \circ \alpha_2$. Now fix a point $t \in U$ in the domain. Then since both f and g are subductive, we can find two plots $\beta_1 : U_1 \rightarrow X_1$ and $\beta_2 : U_2 \rightarrow X_2$, defined on open neighbourhoods of $t \in U$, such that $\alpha_1|_{U_1} = f \circ \beta_1$ and $\alpha_2|_{U_2} = g \circ \beta_2$. Now the plot

$$(\beta_1|_{U_1 \cap U_2}, \beta_2|_{U_1 \cap U_2}) : U_1 \cap U_2 \longrightarrow X_1 \times X_2$$

takes values in the fibred product because

$$r \circ \beta_1|_{U_2} = R \circ f \circ \beta_1|_{U_2} = R \circ \alpha_1|_{U_1 \cap U_2} = L \circ \alpha_2|_{U_1 \cap U_2} = l \circ \beta_2|_{U_1},$$

and we see that it lifts $(\alpha_1, \alpha_2)|_{U_1 \cap U_2}$ along $f \times g$. $\square$

By setting $A = \{*\}$ to be the one-point space, this lemma gives in particular that the product $f \times g$ of two subductions is again a subduction.

To end this section, we should also mention the existence of the notion of a *local subduction* (or *strong subduction*):

Definition 2.24. A smooth surjection $f : X \rightarrow Y$ is called a *local subduction* if for each $y \in Y$, each $x \in f^{-1}(\{y\})$, and any pointed plot of the form $\alpha : (U_\alpha, 0) \rightarrow (Y, y)$, there exists another pointed plot $\beta : (V, 0) \rightarrow (X, x)$, defined on an open neighbourhood $0 \in V \subseteq U_\alpha$, such that $\alpha|_V = f \circ \beta$.

Compare this to a definition of a subduction, where in general the plot β does not have to hit the point x in the domain of f . Note also that *local subduction* does not mean *locally a subduction everywhere*.

Proposition 2.25 ([IZ13a, Article 2.16]). *The local subductions between smooth manifolds are exactly the surjective submersions.*

Due to the above proposition, the notion of a local subduction will be of interest when studying Lie groupoids in the framework of diffeological Morita equivalence we develop below. See Section 7.1.

3. Diffeological Groupoids

We assume that the reader is familiar with the definition of a (Lie) groupoid. A textbook reference for that theory is [Mac05]. To fix our notation, we give here an informal description of a set-theoretic groupoid. A *groupoid* consists of two sets: G_0 and G , together with five *structure maps*. A groupoid will be denoted $G \rightrightarrows G_0$, or just G . Here G_0 is the set of objects of the groupoid, and G is the set of arrows. The five structure maps are

1. The *source map* $\text{src} : G \rightarrow G_0$,
2. The *target map* $\text{trg} : G \rightarrow G_0$,
3. The *unit map* $\text{id} : G_0 \rightarrow G$, mapping $x \mapsto \text{id}_x$,
4. The *inversion map* $\text{inv} : G \rightarrow G$, mapping $g \mapsto g^{-1}$,
5. And the *composition*:

$$\text{comp} : G \times_{G_0}^{\text{src}, \text{trg}} G \longrightarrow G; \quad (g, h) \mapsto g \circ h.$$

The composition is associative, and the identities and inverses behave as such. We say $G \rightrightarrows G_0$ is a *Lie groupoid* if both G and G_0 are smooth manifolds such that the source and target maps are submersions, and each of the other structure maps are smooth. The definition of a diffeological groupoid is a straightforward generalisation of this:

Definition 3.1. A diffeological groupoid is a groupoid internal to the category of diffeological spaces. Concretely, this means that it is a groupoid $G \rightrightarrows G_0$ such that the object space G_0 and arrow space G are endowed with diffeologies that make all of the structure maps smooth.

As diffeology subsumes smooth manifolds, so do diffeological groupoids capture Lie groupoids. Note the main difference with the definition of a Lie groupoid is that we put no extra assumptions on the source and target maps, whereas to make sense of the composition in Lie groupoids we need $G \times_{G_0}^{\text{src}, \text{trg}} G$ to be a smooth manifold, for which it suffices to assume the source and target maps are submersions. We do, however, have:

Proposition 3.2. The source and target maps of a diffeological groupoid are subductions.

Proof. The smooth structure map $u : G_0 \rightarrow G$, sending each object to its identity arrow, is a global smooth section of the source map, and hence by Lemma 2.20(2) the source map must be a subduction. Since the inversion map is a diffeomorphism, it follows that the target map is a subduction as well. $\square$

Definition 3.3. Let $G \rightrightarrows G_0$ be a diffeological groupoid. The isotropy group at $x \in G_0$ is the collection G_x consisting of all arrows in G from and to x :

$$G_x := \text{Hom}_G(x, x) = \text{src}^{-1}(\{x\}) \cap \text{trg}^{-1}(\{x\}).$$

Definition 3.4. Let $G \rightrightarrows G_0$ be a diffeological groupoid. The orbit of an object $x \in G_0$ is defined as

$$\text{Orb}_G(x) := \{y \in G_0 : \exists x \xrightarrow{g} y\} = \text{trg}(\text{src}^{-1}(\{x\})).$$

The orbit space of the groupoid is the space G_0/G consisting of these orbits. We furnish the orbit space with the quotient diffeology from Definition 2.11, so that $\text{Orb}_G : G_0 \rightarrow G_0/G$ is a subduction.

The orbit space of a Lie groupoid is not necessarily (canonically) a smooth manifold. The flexibility of diffeology allows us to study the smooth structure of orbit spaces of all diffeological groupoids. Below we give some examples of diffeological groupoids.

Example 3.5. Let X be a diffeological space, and let R be an equivalence relation on X . We define the *relation groupoid* $X \times_R X \rightrightarrows X$ as follows. The space of arrows consists of exactly those pairs $(x, y) \in X \times X$ such that xRy . With the composition $(z, y) \circ (y, x) := (z, x)$, this becomes a diffeological groupoid. The orbit space $X/(X \times_R X)$ is just the quotient X/R . When X is a smooth manifold, the relation groupoid becomes a Lie groupoid (even when the quotient is not a smooth manifold).

Example 3.6. Let $G \rightrightarrows G_0$ be a diffeological groupoid. We can then consider the subgroupoid of G that only consists of elements in isotropy groups:

$$I_G := \bigcup_{x \in G_0} G_x \subseteq G.$$

This becomes a diffeological groupoid $I_G \rightrightarrows G_0$ called the *isotropy groupoid*. This has been studied in [Bos07, Example 2.1.9] in the context of Lie groupoids. Note that if $G \rightrightarrows G_0$ is a Lie groupoid, then generally I_G is not a submanifold of G , so the isotropy groupoid may no longer be a Lie groupoid.

Example 3.7. The *thin fundamental groupoid* (or *path groupoid*) $\Pi^{\text{thin}}(M)$ of any smooth manifold M is a diffeological groupoid [CLW16, Proposition A.25].

Example 3.8. The *groupoid of Σ -evolutions* of a Cauchy surface is a diffeological groupoid [Gł19, Section II.2.2].

Example 3.9. For any smooth surjection $\pi : X \rightarrow B$ between diffeological spaces, the fibres $X_b := \pi^{-1}(\{b\})$ get the subset diffeology from X . We then have a diffeological groupoid $\mathbf{G}(\pi) \rightrightarrows B$ called the *structure groupoid*, whose space of arrows is defined as

$$\mathbf{G}(\pi) := \bigcup_{a, b \in B} \text{Diff}(X_a, X_b).$$

Structure groupoids play an important rôle in the theory of diffeological fibre bundles [IZ13a, Chapter 8]. In general, they are too big to be Lie groupoids. They also generalise the notion of a *frame groupoid* for a smooth vector bundle. Related to this, in [vdS20, Section 3.4] structure groupoids are used to define a notion of *smooth linear representations* for diffeological groupoids.

Example 3.10. If we are given a diffeological space X , the *germ groupoid* $\text{Germ}(X) \rightrightarrows X$ consists of all *germs* of local diffeomorphisms on X . Even if X itself is a smooth manifold, this is generally not a Lie groupoid. Germ groupoids are used in [IZL18; IZP20]. A detailed construction of the diffeological structure of this groupoid appears in [vdS20, Section 6.1].

4. Diffeological Groupoid Actions and -Bundles

In the following two sections we generalise the theory of Lie groupoid bibundles to the diffeological setting. The development we present here (as in [vdS20, Chapter IV]) is analogous to the development of the Lie version, save that we need to find a suitable replacement for the notion of a surjective submersion. Some of the proofs from the Lie theory can be performed almost *verbatim* in our setting. These proofs already appear in the literature in various places: [Blo08; dHo12; Lan01a; MM05], and also in the different setting of [MZ15]. We adopt many definitions and proofs from those sources, and point out how the diffeological theory subtly differs from the Lie theory. This difference mainly stems from the existence of quotients and fibred products of diffeological spaces, whereas in the Lie theory more care has to be taken. Ultimately, this extra care is what leads to a restricted Morita theorem for Lie groupoids, whereas the diffeological theorem is more general. In this section specifically we introduce *diffeological groupoid actions* and *-bundles*, two notions that form the ingredients for the main theory on bibundles.

4.1 Diffeological groupoid actions

The most basic notion for the upcoming theory is that of a *groupoid action*. For diffeological groupoids, the definition is the same as for Lie groupoids:

Definition 4.1. Take a diffeological groupoid $G \rightrightarrows G_0$, and a diffeological space X . A smooth left groupoid action of G on X along a smooth map $l_X : X \rightarrow G_0$ is a smooth function

$$G \times_{G_0}^{\text{src}, l_X} X \longrightarrow X; \quad (g, x) \longmapsto g \cdot x,$$

satisfying the following three conditions:

1. For $g \in G$ and $x \in X$ such that $\text{src}(g) = l_X(x)$ we have $l_X(g \cdot x) = \text{trg}(g)$.
2. For every $x \in X$ we have $\text{id}_{l_X(x)} \cdot x = x$.
3. We have $h \cdot (g \cdot x) = (h \circ g) \cdot x$ whenever defined, i.e. when $\text{src}(g) = l_X(x)$ and the arrows are composable.

The smooth map $l_X : X \rightarrow G_0$ is called the left moment map. In-line, we denote an action by $G \curvearrowright^{l_X} X$. To save space, we may write $(g, x) \mapsto gx$ instead.

Right actions are defined similarly: a smooth right groupoid action of G on X along $r_X : X \rightarrow G_0$ is a smooth map

$$X \times_{G_0}^{r_X, \text{trg}} G \longrightarrow X; \quad (x, g) \longmapsto xg,$$

satisfying $r_X(xg) = \text{src}(g)$, $x \cdot \text{id}_{r_X(x)} = x$ and $(x \cdot g) \cdot h = x \cdot (g \circ h)$ whenever defined. Note how the rôle of the source and target maps are switched with respect to the definition of a left action. Right actions will be denoted by $X \curvearrowright^{r_X} G$, and r_X is called the right moment map.

Example 4.2. Any diffeological groupoid $G \rightrightarrows G_0$ acts on its own arrow space from the left and right by composition, which gives actions $G \curvearrowright^{\text{trg}} G$ and $G \curvearrowright^{\text{src}} G$ that are both defined by $(g, h) \mapsto g \circ h$.

Definition 4.3. The orbit of a point $x \in X$ in the space of an action $G \curvearrowright^{l_X} X$ is defined as

$$\text{Orb}_G(x) := \{gx : g \in \text{src}^{-1}(\{l_X(x)\})\}.$$

The quotient space (or orbit space) of the action is defined as the collection of all orbits, and denoted X/G . With the quotient diffeology, the orbit projection map $\text{Orb}_G : X \rightarrow X/G$ becomes a subduction.

The following gives a notion of morphism between actions:

Definition 4.4. Consider two smooth groupoid actions $G \curvearrowright^{l_X} X$ and $G \curvearrowright^{l_Y} Y$. A smooth map $\varphi : X \rightarrow Y$ is called G -equivariant if $l_X = l_Y \circ \varphi$ and it commutes with the actions whenever defined: $\varphi(gx) = g\varphi(x)$.

Definition 4.5. The (smooth left) action category $\mathbf{Act}(G \rightrightarrows G_0)$ of a diffeological groupoid $G \rightrightarrows G_0$ is the category consisting of smooth left actions $G \curvearrowright^{l_X} X$ as objects, and G -equivariant maps as morphisms. This forms the analogue of the category of (left) modules from ring theory. We show in Section 6.3 that the action category is in some sense a Morita invariant.

4.1.1 The balanced tensor product

We now give an important construction that will later allow us to define the *composition* of bibundles.

Construction 4.6. Consider a diffeological groupoid $H \rightrightarrows H_0$, with a smooth left action $H \curvearrowright^{l_Y} Y$ and a smooth right action $X \curvearrowright^{r_X} H$. On the fibred product $X \times_{H_0}^{r_X, l_Y} Y$ we define the following smooth right H -action. The moment map is $R := r_X \circ \text{pr}_1|_{X \times_{H_0} Y} = l_Y \circ \text{pr}_2|_{X \times_{H_0} Y}$, and the action is given by:

$$\left(X \times_{H_0}^{r_X, l_Y} Y \right) \times_{H_0}^{R, \text{trg}} H \longrightarrow X \times_{H_0}^{r_X, l_Y} Y; \quad ((x, y), h) \longmapsto (x \cdot h, h^{-1} \cdot y).$$

It is clear that this action is also smooth, and we call it the *diagonal H -action*. The *balanced tensor product* is the diffeological space defined as the orbit space of this smooth groupoid action:

$$X \otimes_H Y := \left(X \times_{H_0}^{r_X, l_Y} Y \right) / H.$$

The orbit of a pair of points (x, y) in the balanced tensor product will be denoted $x \otimes y$. Whenever we encounter a term of the form $x \otimes y \in X \otimes_H Y$, we assume that it is well defined, i.e. $r_X(x) = l_Y(y)$. The terminology is explained by the following useful identity:

$$xh \otimes y = x \otimes hy.$$

In the literature on Lie groupoids, this space is sometimes called the *Hilsum-Skandalis tensor product*, named after a construction appearing in [HS87].

We note that this marks the first difference with the development of the Lie theory of bibundles and Morita equivalence. There, the balanced tensor product can only be defined when both $X \times_{H_0}^{r_X, l_Y} Y$ and the quotient by the diagonal H -action are smooth manifolds. This is usually only done after (bi)bundles are defined, and some principality conditions are presupposed. The principality then exactly ensures the existence of canonical differentiable structures on the fibred product and quotient. Here, the flexibility of diffeology allows us to define the balanced tensor product in an earlier stage of the development, and we do so to demonstrate this conceptual difference.

4.2 Diffeological groupoid bundles

A groupoid bundle is a smooth map, whose domain carries a groupoid action, such that the fibres of the map are preserved by this action:

Definition 4.7. A smooth left diffeological groupoid bundle is a smooth left groupoid action $G \curvearrowright^l X$ together with a G -invariant smooth map $\pi : X \rightarrow B$. We denote such bundles by $G \curvearrowright^l X \xrightarrow{\pi} B$, and also call them (left) G -bundles. Right bundles are defined similarly, and denoted $B \xleftarrow{\pi} X \curvearrowright^r G$.

The next definition gives a notion of morphism between bundles:

Definition 4.8. Take two left G -bundles $G \curvearrowright^l X \xrightarrow{\pi_X} B$ and $G \curvearrowright^l Y \xrightarrow{\pi_Y} B$ over the same base. A G -bundle morphism is a G -equivariant smooth map $\varphi : X \rightarrow Y$ such that $\pi_X = \pi_Y \circ \varphi$. We make a similar definition for right bundles.

In order to define Morita equivalence, we need to define a notion of when a bundle is *principal*. For Lie groupoid bundles, these generalise the ordinary notion of smooth principal bundles of Lie groups and manifolds. That definition involves the notion of a surjective submersion. As we have mentioned, this notion needs to be generalised to diffeology. Proposition 2.25 suggests that we could take *local subductions*, since they directly generalise the surjective submersions. However, it turns out that *subductions* behave sufficiently like submersions for the theory to work. The following definition then generalises the fact that the underlying bundle of a principal Lie groupoid bundle has to be a submersion:

Definition 4.9. A diffeological groupoid bundle $G \curvearrowright^l X \xrightarrow{\pi} B$ is called *subductive* if the underlying map $\pi : X \rightarrow B$ is a subduction.

The following generalises the fact that the action of a principal Lie groupoid bundle has to be free and transitive on the fibres:

Definition 4.10. A diffeological groupoid bundle $G \curvearrowright^l X \xrightarrow{\pi} B$ is called *pre-principal* if the action map $A_G : G \times_{G_0}^{\text{src}, l_X} X \rightarrow X \times_B^{\pi, \pi} X$ mapping $(g, x) \mapsto (gx, x)$ is a diffeomorphism.

Combining these two:

Definition 4.11. *A diffeological groupoid bundle is called principal if it is both subductive and pre-principal.*

This definition serves as our generalisation of principal Lie groupoid bundles, cf. [Blo08, Definition 2.10] and [dHo12, Section 3.6]. Clearly any principal Lie groupoid bundle in the sense described in those references is also a principal diffeological groupoid bundle. Note that in the Lie theory, most constructions (such as the balanced tensor product) depend on the submersiveness of the underlying bundle map, so it makes little sense to define pre-principality for Lie groupoids. However, as we have already seen, in the diffeological case these constructions can be carried out more generally, and this will allow us to see what parts of the development of the theory depend on either the subductiveness or pre-principality of the bundles, rather than on full principality. In our development of the theory, some proofs can therefore be performed separately, whereas in the Lie theory they have to be performed at once. We hope this makes for clearer exposition.

Note also that when a bundle $G \curvearrowright^{\text{lx}} X \xrightarrow{\pi} B$ is pre-principal, the action map induces a diffeomorphism $X/\pi \cong X/G$, and when the bundle is subductive, Example 2.19 gives a diffeomorphism $B \cong X/\pi$. For a principal bundle we therefore have $B \cong X/G$.

Example 4.12. The action of any diffeological groupoid $G \rightrightarrows G_0$ on its own arrow space (Example 4.2) forms a bundle $G \curvearrowright^{\text{trg}} G \xrightarrow{\text{src}} G_0$. From Proposition 3.2 it follows that this bundle is principal.

4.2.1 The division map of a pre-principal bundle

The material in this section is similar to [Blo08, Section 3.1] for Lie groupoids. If a bundle $G \curvearrowright^{\text{lx}} X \xrightarrow{\pi} B$ is pre-principal, the fact that the action map is bijective gives that the action $G \curvearrowright^{\text{lx}} X$ has to be *free*, and *transitive* on the π -fibres. This means that for every two points $x, y \in X$ such that $\pi(x) = \pi(y)$, there exists a *unique* arrow $g \in G$ such that $gy = x$. We denote this arrow by $\langle x, y \rangle_G$, and the map $\langle \cdot, \cdot \rangle_G$ is called the *division map*⁴:

⁴The notational resemblance to an inner-product is not accidental. The division map plays a very similar rôle to the inner product of a Hilbert C^* -module. For more on this analogy, see [Blo08, Section 3].

Definition 4.13. Let $G \curvearrowright^{l_X} X \xrightarrow{\pi} B$ be a pre-principal G -bundle, and let A_G denote its action map. Then the division map associated to this bundle is the smooth map

$$\langle \cdot, \cdot \rangle_G : X \times_B^{\pi, \pi} X \xrightarrow{A_G^{-1}} G \times_{G_0}^{\text{src}, l_X} X \xrightarrow{\text{pr}_1|_{G \times_{G_0} X}} G.$$

We summarise some algebraic properties of the division map that will be used in our proofs throughout later sections. The proofs are straightforward, and use the uniqueness property described above.

Proposition 4.14. Let $G \curvearrowright^{l_X} X \xrightarrow{\pi} B$ be a pre-principal G -bundle. Its division map $\langle \cdot, \cdot \rangle_G$ satisfies the following properties:

1. The source and targets are $\text{src}(\langle x_1, x_2 \rangle_G) = l_X(x_2)$ and $\text{trg}(\langle x_1, x_2 \rangle_G) = l_X(x_1)$.
2. The inverses are given by $\langle x_1, x_2 \rangle_G^{-1} = \langle x_2, x_1 \rangle_G$.
3. For every $x \in X$ we have $\langle x, x \rangle_G = \text{id}_{l_X(x)}$.
4. Whenever well-defined, we have $\langle gx_1, x_2 \rangle_G = g \circ \langle x_1, x_2 \rangle_G$.

Proposition 4.15. Let $\varphi : X \rightarrow Y$ be a bundle morphism between two pre-principal G -bundles $G \curvearrowright^{l_X} X \xrightarrow{\pi_X} B$ and $G \curvearrowright^{l_Y} Y \xrightarrow{\pi_Y} B$. Denoting the division maps of these bundles respectively by $\langle \cdot, \cdot \rangle_G^X$ and $\langle \cdot, \cdot \rangle_G^Y$, we have for all $x_1, x_2 \in X$ in the same π_X -fibre that:

$$\langle x_1, x_2 \rangle_G^X = \langle \varphi(x_1), \varphi(x_2) \rangle_G^Y.$$

Proof. Observe that $\langle \varphi(x_1), \varphi(x_2) \rangle_G^Y$ is the unique arrow that satisfies $\langle \varphi(x_1), \varphi(x_2) \rangle_G^Y \varphi(x_2) = \varphi(x_1)$. However, by G -equivariance we get $\varphi(x_1) = \varphi(\langle x_1, x_2 \rangle_G^X x_2) = \langle x_1, x_2 \rangle_G^X \varphi(x_2)$, from which the claim immediately follows. $\square$

4.2.2 Invertibility of G -bundle morphisms

We now prove a result that generalises the fact that morphisms between principal Lie group bundles are always diffeomorphisms. In our case we shall do the proof in two separate lemmas.

Lemma 4.16. *Consider a G -bundle morphism $\varphi : X \rightarrow Y$ between a pre-principal bundle $G \curvearrowright^{l_X} X \xrightarrow{\pi_X} B$ and a bundle $G \curvearrowright^{l_Y} Y \xrightarrow{\pi_Y} B$ whose underlying action $G \curvearrowright^{l_Y} Y$ is free. Then φ is injective.*

Proof. Since $G \curvearrowright^{l_X} X \xrightarrow{\pi_X} B$ is pre-principal, we get a smooth division map $\langle \cdot, \cdot \rangle_G^X$. To start the proof, suppose that we have two points $x_1, x_2 \in X$ satisfying $\varphi(x_1) = \varphi(x_2)$. Since φ preserves the fibres, we get that

$$\pi_X(x_1) = \pi_Y \circ \varphi(x_1) = \pi_Y \circ \varphi(x_2) = \pi_X(x_2).$$

Hence the pair (x_1, x_2) defines an element in $X \times_B X$, so we get an arrow $\langle x_1, x_2 \rangle_G^X \in G$, satisfying $\langle x_1, x_2 \rangle_G^X x_2 = x_1$. If we apply φ to this equation and use its G -equivariance, we get $\varphi(x_1) = \langle x_1, x_2 \rangle_G^X \varphi(x_2)$. However, by assumption, $\varphi(x_1) = \varphi(x_2)$ and the action $G \curvearrowright^{l_Y} Y$ is free, so we must have that $\langle x_1, x_2 \rangle_G^X$ is the identity arrow at $l_Y \circ \varphi(x_2) = l_X(x_2)$. Hence we get the desired result:

$$x_1 = \langle x_1, x_2 \rangle_G^X x_2 = \text{id}_{l_X(x_2)} x_2 = x_2. \quad \square$$

Lemma 4.17. *Consider a G -bundle morphism $\varphi : X \rightarrow Y$ from a subductive bundle $G \curvearrowright^{l_X} X \xrightarrow{\pi_X} B$ to a pre-principal bundle $G \curvearrowright^{l_Y} Y \xrightarrow{\pi_Y} B$. Then φ is a subduction.*

Proof. Denote the smooth division map of $G \curvearrowright^{l_Y} Y \xrightarrow{\pi_Y} B$ by $\langle \cdot, \cdot \rangle_G^Y$. Then φ and $\langle \cdot, \cdot \rangle_G^Y$ combine into a smooth map

$$\psi : X \times_B^{\pi_X, \pi_Y} Y \longrightarrow X; \quad (x, y) \longmapsto \langle y, \varphi(x) \rangle_G^Y x.$$

Note that this is well defined because if $\pi_X(x) = \pi_Y(y)$, then $\pi_Y \circ \varphi(x) = \pi_Y(y)$ as well, and moreover $l_Y \circ \varphi(x) = l_X(x)$, showing that the action on the right hand side is allowed. The G -equivariance of φ then gives

$$\varphi \circ \psi = \text{pr}_2|_{X \times_B Y}.$$

Since π_X is a subduction, so is $\text{pr}_2|_{X \times_B Y}$ by Lemma 2.22, and by Lemma 2.20(2) it follows φ is a subduction. $\square$

Proposition 4.18. *Any bundle morphism from a principal groupoid bundle to a pre-principal groupoid bundle is a diffeomorphism. In particular, both must then be principal.*

Proof. By Lemma 4.17 any such bundle morphism is a subduction, and since in particular the underlying action of a pre-principal bundle is free, it must also be injective by Lemma 4.16. The result follows by Proposition 2.21. That the second bundle is principal too follows from the fact that a bundle map preserves the fibres, so the projection of the second bundle can be written as the composition of a diffeomorphism and a subduction. $\square$

5. Diffeological Bibundles and Morita Equivalence

This section contains the main definition of this paper: the notion of a *bi-principal bibundle*, which immediately gives our definition of *Morita equivalence*. The definition of groupoid bibundles for diffeology are a straightforward adaptation of the definition in the Lie case:

Definition 5.1. Let $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ be two diffeological groupoids. A diffeological (G, H) -bibundle consists of a smooth left action $G \curvearrowright^l X$ and a smooth right action $X \curvearrowright^r H$ such that the left moment map l_X is H -invariant and the right moment map r_X is G -invariant, and moreover such that the actions commute: $(g \cdot x) \cdot h = g \cdot (x \cdot h)$, whenever defined. We draw:

$$\begin{array}{ccccc} G & \curvearrowright & X & \curvearrowright & H \\ \Downarrow & \swarrow l_X & & \searrow r_X & \Downarrow \\ G_0 & & & & H_0 \end{array}$$

and denote them by $G \curvearrowright^l X \curvearrowright^r H$ in-line. Underlying each bibundle are two groupoid bundles: the left underlying G -bundle $G \curvearrowright^l X \xrightarrow{r_X} H_0$ and the right underlying H -bundle $G_0 \xleftarrow{l_X} X \curvearrowright^r H$. It is the properties of these underlying bundles that will determine the behaviour of the bundle itself.

Definition 5.2. Consider a diffeological bibundle $G \curvearrowright^l X \curvearrowright^r H$. We say this bibundle is left pre-principal if the left underlying bundle $G \curvearrowright^l X \xrightarrow{r_X} H_0$ is pre-principal. We say it is right pre-principal if the right underlying bundle $G_0 \xleftarrow{l_X} X \curvearrowright^r H$ is pre-principal. We make similar definitions for subductiveness and principality. Notice that, in this convention, if a bibundle

$G \curvearrowright^l X \curvearrowright^r H$ is left subductive, then its right moment map r_X is a subduction (and vice versa)⁵.

We now have the main definition of this theory:

Definition 5.3. A diffeological bibundle is called:

1. pre-biprincipal if it is both left- and right pre-principal⁶;
2. bisubductive if it is both left- and right subductive;
3. biprincipal if it is both left- and right principal.

Two diffeological groupoids G and H are called Morita equivalent if there exists a biprincipal bibundle between them, and in that case we write $G \simeq_{\text{ME}} H$.

Compare this to the original definition [MRW87, Definition 2.1] of equivalence for locally compact Hausdorff groupoids. We will prove in Corollary 5.23 that Morita equivalence forms a genuine equivalence relation.

Example 5.4. Since submersions between manifolds are subductions with respect to the manifold diffeologies, we see that if two *Lie* groupoids $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ are Morita equivalent in the *Lie* sense (e.g. [CM18, Definition 2.15]), then they are Morita equivalent in the *diffeological* sense. We remark on the converse question in Section 7.1.

In fact, many elementary examples of Morita equivalences between Lie groupoids generalise straightforwardly to analogously defined diffeological groupoids. We refer to [vdS20, Section 4.3] for some of these examples. For us, the most important one is:

Example 5.5. Consider a diffeological groupoid $G \rightrightarrows G_0$. There exists a canonical (G, G) -bibundle structure on the space of arrows G , which is called the *identity bibundle*. The actions are just the composition in G itself,

⁵Note: [dHo12, Section 4.6] defines this differently, where “[a] bundle is left (resp. right) principal if only the right (resp. left) underlying bundle is so.” We suspect this may be a typo, since it apparently conflicts with their use of terminology in the proof of [dHo12, Theorem 4.6.3]. We stick to the terminology defined above, where *left* principality pertains to the *left* underlying bundle.

⁶The prefixes *bi*- and *pre*- commute: “bi-(pre-principal) = pre-(biprincipal)”.

as in Example 4.2. Note that the identity bibundle is always biprincipal, because the action map has a smooth inverse $(g_1, g_2) \mapsto (g_1 \circ g_2^{-1}, g_2)$. This proves that any diffeological groupoid is Morita equivalent to itself, through the identity bibundle $G \curvearrowright^{\text{trg}} G^{\text{src}} \curvearrowleft G$.

Construction 5.6. Consider a diffeological bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$. The *opposite bibundle* $H \curvearrowright^{l_{\bar{X}}} \bar{X} \curvearrowright^{r_{\bar{X}}} G$ is defined as follows. The underlying diffeological space does not change, $\bar{X} := X$, but the moment maps switch, meaning that $l_{\bar{X}} := r_X$ and $r_{\bar{X}} := l_X$, and the actions are defined as follows:

$$\begin{aligned} H \curvearrowright^{r_X} \bar{X}; \quad & h \cdot x := xh^{-1}, \\ \bar{X} \curvearrowright^{l_X} G; \quad & x \cdot g := g^{-1}x. \end{aligned}$$

Here the actions on the right-hand sides are the original actions of the bibundle. It is easy to see that performing this operation twice gives the original bibundle back. It is also important to note that for all properties defined in Definition 5.2, taking the opposite merely switches the words ‘left’ and ‘right’.

The following extends Proposition 4.14(4):

Lemma 5.7. *Consider a left pre-principal bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$, and also the opposite G -action $\bar{X} \curvearrowright^{l_X} G$. Then, whenever defined, we have:*

$$\langle x_1, x_2 g \rangle_G = \langle x_1, x_2 \rangle_G \circ g.$$

Proof. This follows directly from Proposition 4.14 and the definition of the opposite action:

$$\langle x_1, x_2 g \rangle_G = \langle x_1, g^{-1} x_2 \rangle_G = (g^{-1} \circ \langle x_2, x_1 \rangle_G)^{-1} = \langle x_1, x_2 \rangle_G \circ g. \quad \square$$

5.1 Induced actions

A bibundle $G \curvearrowright X \curvearrowleft H$ allows us to transfer a groupoid action $H \curvearrowright Y$ to a groupoid action $G \curvearrowright X \otimes_H Y$. This is called the *induced action*, and, together with the balanced tensor product, will be crucial to define the composition of bibundles. The idea is that G acts on the first component of $X \otimes_H Y$.

Construction 5.8. Consider a diffeological bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$, and a smooth action $H \curvearrowright^{l_Y} Y$. We construct a smooth left G -action on the balanced tensor product $X \otimes_H Y$. The left moment map is defined as

$$L_X : X \otimes_H Y \longrightarrow G_0; \quad x \otimes y \longmapsto l_X(x).$$

This is well defined because l_X is H -invariant, and smooth by Lemma 2.20(3). For an arrow $g \in G$ with $\text{src}(g) = L_X(x \otimes y) = l_X(x)$, define the action as:

$$G \curvearrowright^{L_X} X \otimes_H Y; \quad g \cdot (x \otimes y) := (gx) \otimes y.$$

Note that the right hand side is well defined because r_X is G -invariant and the G - and H -actions commute, so $r_X(gx) = l_Y(y)$ and the expression does not change if we replace $x \otimes y$ by $xh \otimes h^{-1}y$ for arbitrary $h \in H$. Since there can be no confusion, we will drop all parentheses and write $gx \otimes y$ instead. That the action is smooth follows because $(g, (x, y)) \mapsto (gx, y)$ is smooth (on the appropriate domains) and by another application of Lemma 2.20(3). Hence we obtain the *induced action* $G \curvearrowright^{L_X} X \otimes_H Y$.

Now suppose that we are given a smooth H -equivariant map $\varphi : Y_1 \rightarrow Y_2$ between two smooth actions $H \curvearrowright^{l_1} Y_1$ and $H \curvearrowright^{l_2} Y_2$. We define a map

$$\text{id}_X \otimes \varphi : X \otimes_H Y_1 \longrightarrow X \otimes_H Y_2; \quad x \otimes y \longmapsto x \otimes \varphi(y).$$

The underlying map $X \times_{H_0} Y_1 \rightarrow X \times_{H_0} Y_2 : (x, y) \mapsto (x, \varphi(y))$ is clearly smooth. Then by composition of the projection onto $X \otimes_H Y_2$ and Lemma 2.20(3), we find $\text{id}_X \otimes \varphi$ is smooth. Moreover, it is G -equivariant:

$$\text{id}_X \otimes \varphi(gx \otimes y) = gx \otimes \varphi(y) = g(\text{id}_X \otimes \varphi(x \otimes y)).$$

Definition 5.9. A diffeological bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ defines an induced action functor:

$$\begin{aligned} X \otimes_H - : \mathbf{Act}(H \rightrightarrows H_0) &\longrightarrow \mathbf{Act}(G \rightrightarrows G_0), \\ (H \curvearrowright^{l_Y} Y) &\longmapsto (G \curvearrowright^{L_X} X \otimes_H Y), \\ \varphi &\longmapsto \text{id}_X \otimes \varphi. \end{aligned}$$

sending each smooth left H -action $(H \curvearrowright^{l_Y} Y) \mapsto (G \curvearrowright^{L_X} X \otimes_H Y)$ and each H -invariant map $\varphi \mapsto \text{id}_X \otimes \varphi$. We will use this functor in Section 6.3.

5.2 The bicategory of diffeological groupoids and -bibundles

Combining the balanced tensor product (Construction 4.6) and the induced action of a bibundle (Construction 5.8), we can define a notion of composition for diffeological bibundles, and thereby obtain a new sort of category of diffeological groupoids⁷. Specifically, in Theorem 5.17 we will see that we obtain a *bicategory* **DiffeolBiBund**. A bicategory is like a category where the axioms of composition hold merely up to *canonical 2-isomorphism*. For the precise definition we refer to e.g. [Mac71; Lac10]. The point of this section is to give precise definitions for this bicategorical structure, with the first ingredient being the following:

Definition 5.10. Let $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ and $G \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} H$ be two bibundles between the same two diffeological groupoids. A smooth map $\varphi : X \rightarrow Y$ is called a *bibundle morphism* if it is a bundle morphism between both underlying bundles. We also say that φ is *biequivariant*. Concretely, this means that the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{r_X} & H_0 \\ l_X \downarrow & \searrow \varphi & \uparrow r_Y \\ G_0 & \xleftarrow{l_Y} & Y, \end{array} \quad \text{that is:} \quad \begin{aligned} l_X &= l_Y \circ \varphi, \\ r_X &= r_Y \circ \varphi, \end{aligned}$$

and that φ is equivariant with respect to both actions. These will be the 2-morphisms in **DiffeolBiBund**. The isomorphisms of bibundles are exactly the diffeomorphic biequivariant maps. These will be the 2-isomorphisms in **DiffeolBiBund**.

The composition of bibundles is defined as follows:

Construction 5.11. Consider two diffeological bibundles $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ and $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} K$. We shall define on $X \otimes_H Y$ a (G, K) -bibundle structure using the induced actions from Construction 5.8. On the left we take the induced G -action along $L_X : X \otimes_H Y \rightarrow G_0$, which we recall maps $x \otimes y \mapsto l_X(x)$, defined by

$$G \curvearrowright^{L_X} X \otimes_H Y; \quad g(x \otimes y) := (gx) \otimes y.$$

⁷The most straightforward way to obtain a (2-)category of diffeological groupoids is to consider the *smooth functors* and *smooth natural transformations*. We will not be studying this category in the current paper.

Analogous to Construction 5.8, swapping left actions for right actions, we get an induced K -action on the right along $R_Y : X \otimes_H Y \rightarrow K_0$, which maps $x \otimes y \mapsto r_Y(y)$, given by

$$X \otimes_H Y \overset{R_Y}{\curvearrowright} K; \quad (x \otimes y)k := x \otimes (yk).$$

It is easy to see that these two actions form a bibundle $G \overset{L_X}{\curvearrowright} X \otimes_H Y \overset{R_Y}{\curvearrowright} K$, which we also call the *balanced tensor product*.

From this construction we can see that the composition of bibundles will not be strictly associative, and this is where the bicategorical structure becomes important. The following two propositions characterise the compositional structure of the balanced tensor product *up to biequivariant diffeomorphism*. The first of these shows that the identity bibundle (Example 5.5) is a *weak identity*:

Proposition 5.12. *Let $G \overset{L_X}{\curvearrowright} X \overset{r_X}{\curvearrowright} H$ be a diffeological bibundle. Then there are biequivariant diffeomorphisms*

$$\begin{array}{ccc} G \overset{L_G}{\curvearrowright} G \otimes_G X \overset{R_X}{\curvearrowright} H & & G \overset{L_X}{\curvearrowright} X \otimes_H H \overset{R_H}{\curvearrowright} H \\ \Downarrow \varphi & \text{and} & \Downarrow \\ G \overset{L_X}{\curvearrowright} X \overset{r_X}{\curvearrowright} H & & G \overset{L_X}{\curvearrowright} X \overset{r_X}{\curvearrowright} H. \end{array}$$

Proof. The idea of the proof is briefly sketched on [Blo08, p.8]. The map $\varphi : G \otimes_G X \rightarrow X$ is defined by the action: $g \otimes x \mapsto gx$. This map is clearly well defined, and by an easy application of Lemma 2.20(3) also smooth. Further note that φ intertwines the left moment maps:

$$l_X \circ \varphi(g \otimes x) = l_X(gx) = \text{trg}(g) = L_G(g \otimes x),$$

and similarly we find it intertwines the right moment maps. Associativity of the G -action and the fact that it commutes with the H -action directly ensure that φ is biequivariant. Moreover, we claim that the smooth map $\psi : X \rightarrow G \otimes_G X$ defined by $x \mapsto \text{id}_{l_X(x)} \otimes x$ is the inverse of φ . It follows easily that $\varphi \circ \psi = \text{id}_X$, and the other side follows from the defining property of the balanced tensor product:

$$\psi \circ \varphi(g \otimes x) = \psi(gx) = \text{id}_{l_X(gx)} \otimes gx = (\text{id}_{\text{trg}(g)} \circ g) \otimes x = g \otimes x.$$

It follows from an analogous argument that the identity bibundle of H acts like a weak right inverse. $\square$

The second proposition shows that the balanced tensor product is associative *up to canonical biequivariant diffeomorphism*:

Proposition 5.13. *Let $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$, $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} H'$, and $H' \curvearrowright^{l_Z} Z \curvearrowright^{r_Z} K$ be diffeological bibundles. Then there exists a biequivariant diffeomorphism*

$$\begin{array}{ccc} G \curvearrowright^{L_{X \otimes_H Y}} (X \otimes_H Y) \otimes_{H'} Z \curvearrowright^{R_Z} K & & \\ \Downarrow A & & A : (x \otimes y) \otimes z \longmapsto x \otimes (y \otimes z). \\ G \curvearrowright^{L_X} X \otimes_H (Y \otimes_{H'} Z) \curvearrowright^{R_{Y \otimes_{H'} Z}} K, & & \end{array}$$

Proof. That the map A is smooth follows by Lemma 2.20(3), because the corresponding underlying map $((x, y), z) \mapsto (x, (y, z))$ is a diffeomorphism. The inverse of this diffeomorphism on the underlying fibred product induces exactly the smooth inverse of A , showing that A is a diffeomorphism. Furthermore, it is easy to check that A is biequivariant. $\square$

Combining Propositions 5.12 and 5.13 gives that the balanced tensor product of bibundles does indeed behave like the composition in a bicategory. Next to the composition of arrows in a bicategory, we also need to describe the compositional structure of the 2-arrows. The following elementary result says that the ordinary *vertical composition* of biequivariant maps is again biequivariant:

Proposition 5.14. *Consider two biequivariant smooth maps:*

$$\begin{array}{ccc} G \curvearrowright^{l_X} X \curvearrowright^{r_X} H & & \\ \Downarrow \varphi & & \\ G \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} H & & \\ \Downarrow \psi & & \\ G \curvearrowright^{l_Z} Z \curvearrowright^{r_Z} H. & & \end{array}$$

Then the composition $\psi \circ \varphi : X \rightarrow Z$ is also biequivariant.

Next to vertical composition, a bicategory should also allow for *horizontal composition* of 2-arrows. Again, the construction of this composition follows the Lie groupoid theory:

Construction 5.15. Consider the following situation of four bibundles and two biequivariant maps:

$$\begin{array}{ccc} G \curvearrowright^{l_X} X \curvearrowright^{r_X} H & & H \curvearrowright^{l_P} P \curvearrowright^{r_P} K \\ \downarrow \varphi & \text{and} & \downarrow \psi \\ G \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} H & & H \curvearrowright^{l_Q} Q \curvearrowright^{r_Q} K. \end{array}$$

The goal will be to construct a biequivariant map

$$\begin{array}{c} G \curvearrowright^{L_X} X \otimes_H P \curvearrowright^{R_P} K \\ \downarrow \varphi \otimes \psi \\ G \curvearrowright^{L_Y} Y \otimes_H Q \curvearrowright^{R_Q} K, \end{array}$$

called the *horizontal composition* of φ and ψ . The most obvious choice for the underlying function is the following:

$$\varphi \times \psi|_{X \times_H P} : X \times_H^{r_X, l_P} P \longrightarrow Y \times_H^{r_Y, l_Q} Q.$$

The biequivariance of φ and ψ ensures that the image of this function indeed lands in the fibred product $Y \times_H^{r_Y, l_Q} Q$, showing it is well defined and smooth. Projecting down to the balanced tensor products, we define:

$$\varphi \otimes \psi : X \otimes_H P \longrightarrow Y \otimes_H Q; \quad x \otimes p \longmapsto \varphi(x) \otimes \psi(p).$$

To show that this will again form a 2-arrow in **DiffeolBiBund**, we have the following counterpart to Proposition 5.14:

Proposition 5.16. *The map $\varphi \otimes \psi : X \otimes_H P \rightarrow Y \otimes_H Q$ from Construction 5.15 is a well-defined smooth biequivariant map of diffeological bibundles.*

Proof. We start by showing that $\varphi \otimes \psi$ is a well-defined function on the balanced tensor products. For that, take an element $x \otimes p \in X \otimes_H P$, and take an arbitrary arrow $h \in H$ with $\text{trg}(h) = r_X(x) = l_P(p)$, such that $x \otimes p = xh \otimes h^{-1}p$. Using the fact that both φ and ψ are biequivariant, and

the defining relation of the balanced tensor product $Y \otimes_H Q$, we calculate:

$$\begin{aligned}\varphi \otimes \psi(xh \otimes h^{-1}p) &:= \varphi(xh) \otimes \psi(h^{-1}p) \\ &= \varphi(x)h \otimes h^{-1}\psi(p) \\ &= \varphi(x) \otimes \psi(p) \\ &=: \varphi \otimes \psi(x \otimes p),\end{aligned}$$

showing that $\varphi \otimes \psi$ is indeed well defined. Next, observe that we have the following commutative diagram of functions:

$$\begin{array}{ccc} X \times_H^{r_X, l_P} P & \xrightarrow{\varphi \times \psi|_{X \times_H P}} & Y \times_H^{r_Y, l_Q} Q \\ \downarrow & & \downarrow \\ X \otimes_H P & \xrightarrow{\varphi \otimes \psi} & Y \otimes_H Q, \end{array}$$

where the vertical arrows are the canonical projections. It follows immediately from Lemma 2.20(3) that $\varphi \otimes \psi$ is also smooth.

Lastly, we show that $\varphi \otimes \psi$ is biequivariant with respect to the G - and K -actions. An easy calculation using the biequivariance of φ shows that

$$\begin{aligned}L_Y \circ (\varphi \otimes \psi)(x \otimes p) &= L_Y(\varphi(x) \otimes \psi(p)) \\ &= l_Y \circ \varphi(x) \\ &= l_X(x) \\ &= L_X(x \otimes p),\end{aligned}$$

and similarly we find $R_Q \circ (\varphi \otimes \psi) = R_P$. Moreover, $\varphi \otimes \psi$ commutes with the left G -actions:

$$\begin{aligned}\varphi \otimes \psi(g \cdot (x \otimes p)) &= \varphi \otimes \psi((gx) \otimes p) \\ &= \varphi(gx) \otimes \psi(p) \\ &= (g\varphi(x)) \otimes \psi(p) \\ &= g \cdot (\varphi(x) \otimes \psi(p)) \\ &= g \cdot (\varphi \otimes \psi(x \otimes p)),\end{aligned}$$

and we similarly find that it commutes with the right K -actions. We have thus proved that $\varphi \otimes \psi$ defines a smooth biequivariant map, as desired. $\square$

We have now described all bicategorical ingredients for **DiffeolBiBund**, and so it remains to check that they do indeed satisfy the axioms of a bicategory. The proof of this is directly analogous to the one for the Lie theory, which is explained in [Blo08, Proposition 2.12], and we therefore leave out the details.

Theorem 5.17. *There is a bicategory **DiffeolBiBund** consisting of diffeological groupoids as objects, diffeological bibundles as morphisms with balanced tensor product as composition, and biequivariant smooth maps as 2-morphisms.*

Proof idea. Note that the way in which we have defined the bicategorical structure of **DiffeolBiBund** is a direct generalisation of the Lie groupoid theory (in the sense that, when restricted to Lie groupoids, it is the exact same). Furthermore, that the axioms of a bicategory hold for Lie groupoids ([Blo08, Proposition 2.12]) is not dependent on the (left or right) principality of the bibundles (save for the fact that this is needed to ensure the existence of the balanced tensor product), but is rather a property of the underlying functions. Given the results in this section, it is therefore clear that those proofs generalise directly to the diffeological setting. $\square$

As we remarked in Section 4.1, the balanced tensor product for Lie groupoids can only be constructed for *left* (or *right*) *principal* bibundles. This means that in the Lie theory, the category of bibundles only consists of the left (or right) principal bibundles, since otherwise the composition cannot be defined. For diffeology we obtain a bicategory of *all* bibundles.

5.3 Properties of bibundles under composition and isomorphism

We study how the properties of diffeological bibundles defined in Definition 5.2 are preserved under the balanced tensor product and biequivariant diffeomorphism. These results will be crucial in characterising the weakly invertible bibundles. First we show that left subductive and left pre-principal bibundles are closed under composition.

Proposition 5.18. *The balanced tensor product preserves left subductiveness.*

Proof. Consider the balanced tensor product $G \curvearrowright^{L_X} X \otimes_H Y \curvearrowright^{R_Y} K$ of two left subductive bibundles $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ and $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} K$. We need to show that the right moment map $R_Y : X \otimes_H Y \rightarrow K_0$ is a subduction. But, note that it fits into the following commutative diagram:

$$\begin{array}{ccc} X \times_{H_0}^{r_X, l_Y} Y & \xrightarrow{\pi} & X \otimes_H Y \\ \text{pr}_2|_{X \times_{H_0} Y} \downarrow & & \downarrow R_Y \\ Y & \xrightarrow{r_Y} & K_0. \end{array}$$

Here π is the canonical quotient projection. The restricted projection $\text{pr}_2|_{X \times_{H_0} Y}$ is a subduction by Lemma 2.22, since r_X is a subduction. Moreover, r_Y is a subduction, so the bottom part of the diagram is a subduction. It follows by Lemma 2.20(3) that R_Y is a subduction. $\square$

Note that, even though R_Y only explicitly depends on the moment map r_Y , the proof still depends on the subductiveness of r_X as well.

To prove that the balanced tensor product of two left pre-principal bibundles is again left pre-principal, we need the following lemma, describing how the division map interacts with the bibundle structure, extending the list in Proposition 4.14 on the algebraic properties of the division map.

Lemma 5.19. *Let $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ be a left pre-principal bibundle, and denote its division map by $\langle \cdot, \cdot \rangle_G$. Then, whenever defined:*

$$\langle x_1, x_2 h \rangle_G = \langle x_1 h^{-1}, x_2 \rangle_G, \quad \text{or equivalently:} \quad \langle x_1 h, x_2 h \rangle_G = \langle x_1, x_2 \rangle_G.$$

Proof. The arrow $\langle x_1 h, x_2 h \rangle_G \in G$ is the unique one that sends $x_2 h$ to $x_1 h$. Now, since the actions commute, we can multiply both of these terms from the right by h^{-1} , which gives the equation $\langle x_1 h, x_2 h \rangle_G x_2 = x_1$, and this immediately gives our result. $\square$

Proposition 5.20. *The balanced tensor product preserves left pre-principality.*

Proof. To start the proof, take two left pre-principal bibundles, with our usual notation: $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ and $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} K$. Denote their division maps by $\langle \cdot, \cdot \rangle_G^X$ and $\langle \cdot, \cdot \rangle_H^Y$, respectively. Using these, we will construct a

smooth inverse of the action map of the balanced tensor product. Let us denote the action map of the balanced tensor product by

$$\Phi : G \times_{G_0}^{\text{src}, L_X} (X \otimes_H Y) \longrightarrow (X \otimes_H Y) \times_{K_0}^{R_Y, R_Y} (X \otimes_H Y),$$

mapping $(g, x \otimes y) \mapsto (gx \otimes y, x \otimes y)$. After some calculations (which we describe below), we propose the following map as an inverse for Φ :

$$\begin{aligned} \Psi : (X \otimes_H Y) \times_{K_0}^{R_Y, R_Y} (X \otimes_H Y) &\longrightarrow G \times_{G_0}^{\text{src}, L_X} (X \otimes_H Y); \\ (x_1 \otimes y_1, x_2 \otimes y_2) &\longmapsto \left(\langle x_1 \langle y_1, y_2 \rangle_H^Y, x_2 \rangle_G^X, x_2 \otimes y_2 \right). \end{aligned}$$

It is straightforward to check that every action and division occurring in this expression is well defined. We need to check that Ψ is independent on the representations of $x_1 \otimes y_1$ and $x_2 \otimes y_2$. Only the first component Ψ_1 of Ψ could be dependent on the representations, so we focus there. Suppose we have two arrows $h_1, h_2 \in H$ satisfying $\text{trg}(h_i) = r_X(x_i) = l_Y(y_i)$, so that $x_i h_i \otimes h_i^{-1} y_i = x_i \otimes y_i$. For the division of y_2 and y_1 we then use Proposition 4.14 to get:

$$\begin{aligned} \langle h_1^{-1} y_1, h_2^{-1} y_2 \rangle_H^Y &= h_1^{-1} \circ \langle y_1, h_2^{-1} y_2 \rangle_H^Y \\ &= h_1^{-1} \circ (h_2^{-1} \circ \langle y_2, y_1 \rangle_H^Y)^{-1} \\ &= h_1^{-1} \circ \langle y_1, y_2 \rangle_H^Y \circ h_2. \end{aligned}$$

Then, using this and Lemma 5.19, we get:

$$\begin{aligned} \Psi_1(x_1 h_1 \otimes h_1^{-1} y_1, x_2 h_2 \otimes h_2^{-1} y_2) &= \langle x_1 h_1 \langle h_1^{-1} y_1, h_2^{-1} y_2 \rangle_H^Y, x_2 h_2 \rangle_G^X \\ &= \langle (x_1 h_1) (h_1^{-1} \circ \langle y_1, y_2 \rangle_H^Y \circ h_2), x_2 h_2 \rangle_G^X \\ &= \langle (x_1 \langle y_1, y_2 \rangle) h_2, x_2 h_2 \rangle_G^X \\ &= \langle x_1 \langle y_1, y_2 \rangle_H^Y, x_2 \rangle_G^X. \end{aligned}$$

Since the second component of Ψ is, by construction, independent on the representation, it follows that Ψ is a well-defined function. We now need to show that Ψ is smooth. The second component is clearly smooth, because it is just the projection onto the second component of the fibred product. That the other component is smooth follows from Lemmas 2.20 and 2.23. Writing

$$\psi : ((x_1, y_1), (x_2, y_2)) \longmapsto \langle x_1 \langle y_1, y_2 \rangle_H^Y, x_2 \rangle_G^X$$

and $\pi : X \times_{H_0}^{r_X, l_Y} Y \rightarrow X \otimes_H Y$ for the canonical projection, we get a commutative diagram

$$\begin{array}{ccc}
 \left(X \times_{H_0}^{r_X, l_Y} Y \right) \times_{K_0}^{\overline{r_Y}, \overline{r_Y}} \left(X \times_{H_0}^{r_X, l_Y} Y \right) & \xrightarrow{\psi} & G. \\
 \downarrow (\pi \times \pi)|_{\text{dom}(\psi)} & & \uparrow \Psi_1 \\
 (X \otimes_H Y) \times_{K_0}^{R_Y, R_Y} (X \otimes_H Y) & &
 \end{array}$$

Here we temporarily use the notation $\overline{r_Y} := r_Y \circ \text{pr}_2|_{X \times_{H_0} Y}$, which satisfies $R_Y \circ \pi = \overline{r_Y}$. Therefore by Lemma 2.23 the left arrow in this diagram is a subduction. Since the map ψ is evidently smooth, it follows by Lemma 2.20(3) that the first component Ψ_1 , and hence Ψ itself, must be smooth.

Thus, we are left to show that Ψ is an inverse for Φ . That Ψ is a right inverse for Φ now follows by simple calculation using Proposition 4.14 and Lemma 5.19:

$$\begin{aligned}
 \Psi \circ \Phi(g, x \otimes y) &= \Psi(gx \otimes y, x \otimes y) \\
 &= (\langle gx \langle y, y \rangle_H^Y, x \rangle_G^X, x \otimes y) \\
 &= (g \circ \langle x, x \rangle_G^X, x \otimes y) \\
 &= (g, x \otimes y).
 \end{aligned}$$

For the other direction, we calculate:

$$\begin{aligned}
 \Phi \circ \Psi(x_1 \otimes y_1, x_2 \otimes y_2) &= \Phi \left(\langle x_1 \langle y_1, y_2 \rangle_H^Y, x_2 \rangle_G^X, x_2 \otimes y_2 \right) \\
 &= \left(\langle x_1 \langle y_1, y_2 \rangle_H^Y, x_2 \rangle_G^X x_2 \otimes y_2, x_2 \otimes y_2 \right) \\
 &= (x_1 \langle y_1, y_2 \rangle_H^Y \otimes y_2, x_2 \otimes y_2) \\
 &= (x_1 \otimes \langle y_1, y_2 \rangle_H^Y y_2, x_2 \otimes y_2) \\
 &= (x_1 \otimes y_1, x_2 \otimes y_2).
 \end{aligned}$$

Here in the second to last step we use the properties of the balanced tensor product to move the arrow $\langle y_1, y_2 \rangle_H^Y$ over the tensor symbol. Hence we conclude that Φ is a diffeomorphism, which proves that $G \curvearrowright^{L_X} X \otimes_H Y \overset{R_Y}{\curvearrowright} K$ is a left pre-principal bibundle. $\square$

Next we show that left subductiveness and left pre-principality are also preserved under biequivariant diffeomorphism.

Proposition 5.21. *Left pre-principality is preserved by biequivariant diffeomorphism.*

Proof. Suppose that $\varphi : X \rightarrow Y$ is a biequivariant diffeomorphism from a left pre-principal bibundle $G \curvearrowright^{l_X} X \overset{r_X}{\curvearrowright} H$ to another diffeological bibundle $G \curvearrowright^{l_Y} Y \overset{r_Y}{\curvearrowright} H$. Denote their left action maps by A_X and A_Y , respectively. The following square commutes because of biequivariance:

$$\begin{array}{ccc} G \times_{G_0}^{\text{src}, l_X} X & \xrightarrow{A_X} & X \times_{H_0}^{r_X, r_X} X \\ (\text{id}_G \times \varphi)|_{G \times_{G_0} X} \downarrow & & \downarrow (\varphi \times \varphi)|_{X \times_{H_0} X} \\ G \times_{G_0}^{\text{src}, l_Y} Y & \xrightarrow{A_Y} & Y \times_{H_0}^{r_Y, r_Y} Y. \end{array}$$

It is easy to see that both vertical maps are diffeomorphisms. Hence it follows A_Y must be a diffeomorphism as well. $\square$

Proposition 5.22. *Left subductiveness is preserved by biequivariant diffeomorphism.*

Proof. Suppose that $\varphi : X \rightarrow Y$ is a biequivariant diffeomorphism from a left subductive bibundle $G \curvearrowright^{l_X} X \overset{r_X}{\curvearrowright} H$ to $G \curvearrowright^{l_Y} Y \overset{r_Y}{\curvearrowright} H$. That the first bundle is left subductive means that r_X is a subduction, but since φ intertwines the moment maps, it follows immediately that $r_Y = r_X \circ \varphi^{-1}$ is a subduction as well. $\square$

Of course, these four propositions all hold for their respective ‘right’ versions as well. This can be proved formally, without repeating the work, by using opposite bibundles.

Corollary 5.23. *Morita equivalence defines an equivalence relation between diffeological groupoids.*

Proof. Morita equivalence is reflexive by the existence of identity bibundles, which are always biprincipal (Example 5.5). It is also easy to check that the opposite bibundle (Construction 5.6) of a biprincipal bibundle is again biprincipal, showing that Morita equivalence is symmetric. Transitivity follows directly from Propositions 5.18 and 5.20 and their opposite versions. $\square$

5.4 Weak invertibility of diffeological bibundles

In this section we prove the main Morita Theorem 5.31. As we explained in the Introduction, in the bicategory of diffeological groupoids we get a notion of *weak isomorphism*. Let us describe these explicitly: a bibundle $G \curvearrowright X \curvearrowright H$ is weakly invertible if and only if there exists a second bibundle $H \curvearrowright Y \curvearrowright G$, such that $X \otimes_H Y$ is biequivariantly diffeomorphic to G and $Y \otimes_G X$ is biequivariantly diffeomorphic to H . The Morita theorem says that such a weak inverse exists if and only if the bibundle is biprincipal. Let us recall the corresponding statement in the Lie theory: a (say) left principal bibundle has a left principal weak inverse if and only if it is biprincipal [Lan01b, Proposition 4.21]. Here both the original bibundle and its weak inverse have to be left principal, since everything takes place in a bicategory of Lie groupoids and left principal bibundles. According to Theorem 5.17 we get a bicategory of arbitrary bibundles, and the question of weak invertibility becomes a slightly more general one, since we do not start out with a bibundle that is already left principal. Instead we have to infer left principality from bare weak invertibility, where neither the weak inverse may be assumed to be left principal.

One direction of the claim in the main theorem is relatively straightforward, and is the same as for Lie groupoids:

Proposition 5.24. *Let $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ be a biprincipal bibundle. Then its opposite bundle $H \curvearrowright^{r_X} \overline{X} \curvearrowright^{l_X} G$ is a weak inverse.*

Proof. We construct biequivariant diffeomorphisms

$$\begin{array}{ccc} G \curvearrowright^{L_X} X \otimes_H \overline{X} \curvearrowright^{R_{\overline{X}}} G & & H \curvearrowright^{L_{\overline{X}}} \overline{X} \otimes_G X \curvearrowright^{R_X} H \\ \varphi_G \Downarrow & \text{and} & \varphi_H \Downarrow \\ G \curvearrowright^{\text{trg}} G^{\text{src}} \curvearrowright G, & & H \curvearrowright^{\text{trg}} H^{\text{src}} \curvearrowright H. \end{array}$$

Since the original bundle is pre-biprincipal, we have a division map $\langle \cdot, \cdot \rangle_G : X \times_{H_0}^{r_X, r_X} \overline{X} \rightarrow G$. We define a new function

$$\varphi_G : X \otimes_H \overline{X} \longrightarrow G; \quad x_1 \otimes x_2 \longmapsto \langle x_1, x_2 \rangle_G.$$

This is independent on the representation of the tensor product by Lemma 5.19, and smooth by Lemma 2.20(3) since $\varphi_G \circ \pi = \langle \cdot, \cdot \rangle_G$, where π is the canonical projection onto the balanced tensor product. We check that φ_G is

biequivariant. It is easy to check that φ_G intertwines the moment maps, for example:

$$\text{src} \circ \varphi_G(x_1 \otimes x_2) = \text{src}(\langle x_1, x_2 \rangle_G) = l_X(x_2) = R_{\overline{X}}(x_1 \otimes x_2).$$

The left G -equivariance of φ_G follows directly out of Proposition 4.14, and the right G -equivariance follows from Lemma 5.7. Hence φ_G is a genuine bibundle morphism.

Since the original bundle is biprincipal, so is its opposite, and therefore by Propositions 5.18 and 5.20 it follows that both balanced tensor products are also biprincipal. Therefore φ_G is in particular a left G -equivariant bundle morphism from a principal bundle $G \curvearrowright^{L_X} X \otimes_H \overline{X} \xrightarrow{R_{\overline{X}}} G_0$ to a pre-principal bundle $G \curvearrowright^{\text{trg}} G \xrightarrow{\text{src}} G_0$, and hence a diffeomorphism by Proposition 4.18. This proves that the opposite bibundle is a weak right inverse. Note that we already need full biprincipality of the original bibundle for this. To prove that it is also a weak left inverse we make an analogous construction for φ_H , which we leave to the reader. $\square$

The rest of this section will be dedicated to proving the converse of this claim, i.e., that a weakly invertible bibundle is biprincipal. First let us remark that by imitating a result from the Lie theory, we can obtain a partial result in this direction. Let us denote by $\mathbf{DiffeolBiBund}_{\text{LP}}$ the bicategory of diffeological groupoids and left principal bibundles. Note that by Section 5.3 left principality is preserved by the balanced tensor product, so this indeed forms a subcategory.

Theorem 5.25. *A left principal diffeological bibundle has a left principal weak inverse if and only if it is biprincipal. That is, the weakly invertible bibundles in $\mathbf{DiffeolBiBund}_{\text{LP}}$ are exactly the biprincipal ones.*

Proof. This follows by combining Proposition 5.24 with an adaptation of an argument from the Lie groupoid theory as in [MM05, Proposition 2.9]. A more detailed proof (for diffeological groupoids) is in [vdS20, Proposition 4.61]. $\square$

This theorem is the most direct analogue of [Lan01b, Proposition 4.21] in the setting of diffeology. Our main theorem will be a further generalisation of this, which says that the same claim holds in the larger bicategory

DiffeolBiBund of *all* bibundles. We break the proof down in several steps, starting with the implication of bisubductiveness:

Proposition 5.26. *A weakly invertible diffeological bibundle is bisubductive.*

Proof. Suppose we have a bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ that admits a weak inverse $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} G$. Let us denote the included biequivariant diffeomorphisms by $\varphi_G : X \otimes_H Y \rightarrow G$ and $\varphi_H : Y \otimes_G X \rightarrow H$, as usual. Since the identity bibundles of G and H are both biprincipal, it follows by Proposition 5.22 that the moment maps L_X , R_X , L_Y and R_Y are all subductions. Together with the original moment maps, we get four commutative squares, each of the form:

$$\begin{array}{ccc} X \times_{H_0}^{r_X, l_Y} Y & \xrightarrow{\pi} & X \otimes_H Y \\ \text{pr}_1|_{X \times_{H_0} Y} \downarrow & & \downarrow L_X \\ X & \xrightarrow{l_X} & G_0. \end{array}$$

Here $\pi : X \times_{H_0}^{r_X, l_Y} Y \rightarrow X \otimes_H Y$ is the quotient map of the diagonal H -action. By Lemma 2.20(3) it follows that, since L_X is a subduction, so is the composition $l_X \circ \text{pr}_1|_{X \times_{H_0} Y}$, and in turn by Lemma 2.20(2) it follows l_X is a subduction. In a similar fashion we find that r_X , l_Y and r_Y are all subductions as well. $\square$

This proposition gets us halfway to proving that weakly invertible bibundles are biprincipal. To prove that they are pre-biprincipal, it is enough to construct smooth division maps. We will give this construction below (Construction 5.29), which follows from a careful reverse engineering of the division map of a pre-principal bundle. Recall from Proposition 5.20 that the smooth inverse of the action map contains the information of both the G -division map and the H -division map. Specifically, the first component of the inverse is of the form $\langle x_1 \langle y_1, y_2 \rangle_H^Y, x_2 \rangle_G^X$, in which if we set $y_1 = y_2$, we simply reobtain the G -division map $\langle x_1, x_2 \rangle_G^X$. The question is if this “reobtaining” can be done in a smooth way. This is not so obvious at first. Namely, if we vary (x_1, x_2) smoothly within $X \times_{H_0}^{r_X, r_X} X$, can we guarantee that y_1 and y_2 vary smoothly with it, while still retaining the equalities $r_X(x_i) = l_Y(y_i)$ and $y_1 = y_2$? The elaborate Construction 5.29 proves that

this can indeed be done. An essential part of our argument will be supplied by the following two lemmas.

Lemma 5.27. *When $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ is a weakly invertible bibundle, admitting a weak inverse $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} G$, then all four actions are free.*

Proof. This follows from an argument that is used in the proof of [Blo08, Proposition 3.23]. Suppose we have an arrow $h \in H$ and a point $y \in Y$ such that $hy = y$. By Proposition 5.26 it follows that in particular l_X is surjective, so we can find $x \in X$ such that $y \otimes x \in Y \otimes_G X$. Then

$$h(y \otimes x) = (hy) \otimes x = y \otimes x.$$

But by Proposition 5.21 the bundle $H \curvearrowright^{l_Y} Y \otimes_G X \xrightarrow{R_X} G_0$, which is equivariantly diffeomorphic to the identity bundle on H , is pre-principal. So, the left action $H \curvearrowright Y \otimes_G X$ is free, and hence $h = \text{id}_{L_Y(y \otimes x)} = \text{id}_{l_Y(y)}$, proving that $H \curvearrowright Y$ is also free. That the three other actions are free follows analogously. $\square$

Lemma 5.28. *Let $X \curvearrowright^{r_X} H$ and $H \curvearrowright^{l_Y} Y$ be smooth actions, so that we can form the balanced tensor product $X \otimes_H Y$. Suppose that $H \curvearrowright Y$ is free. Then $x_1 \otimes y = x_2 \otimes y$ if and only if $x_1 = x_2$. Similarly, if $X \curvearrowright H$ is free, then $x \otimes y_1 = x \otimes y_2$ if and only if $y_1 = y_2$.*

Proof. If $x_1 = x_2$ to begin with, the implication is trivial. Suppose therefore that $x_1 \otimes y = x_2 \otimes y$, which means that there exists an arrow $h \in H$ such that $(x_1 h^{-1}, hy) = (x_2, y)$. In particular $hy = y$, which, because the action on Y is free, implies $h = \text{id}_{l_Y(y)}$, and it follows that $x_1 = x_1 \text{id}_{l_Y(y)}^{-1} = x_2$. $\square$

We shall now describe how the division map arises from local data:

Construction 5.29. For this construction to work, we start with a diffeological bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$, admitting a weak inverse $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} G$. Then consider a pointed plot $\alpha : (U_\alpha, 0) \rightarrow (X \times_{H_0}^{r_X, r_X} X, (x_1, x_2))$. We split α into the components (α_1, α_2) , which in turn give two pointed plots $\alpha_i : (U_\alpha, 0) \rightarrow (X, x_i)$ satisfying $r_X \circ \alpha_1 = r_X \circ \alpha_2 : U_\alpha \rightarrow H_0$. This equation gives a plot of H_0 , and since by Proposition 5.26 the moment map $l_Y : Y \rightarrow H_0$ is a subduction, for every $t \in U_\alpha$ we can find a plot $\beta : V \rightarrow Y$, defined on an open neighbourhood $t \in V \subseteq U_\alpha$, such

that $r_X \circ \alpha_i|_V = l_Y \circ \beta$. From this equation it follows that the smooth maps $(\alpha_i|_V, \beta) : V \rightarrow X \times_{H_0}^{r_X, l_Y} Y$ define two plots of the underlying space of the balanced tensor product. Applying the canonical quotient projection map $\pi : X \times_{H_0}^{r_X, l_Y} Y \rightarrow X \otimes_H Y$, we thus get two full-fledged plots $s \mapsto \alpha_i|_V(s) \otimes \beta(s)$ of the balanced tensor product. We combine these two plots to define yet another smooth map:

$$\Omega^\alpha|_V := (\pi \circ (\alpha_1|_V, \beta), \pi \circ (\alpha_2|_V, \beta)) : V \longrightarrow (X \otimes_H Y) \times_{G_0}^{R_Y, R_Y} (X \otimes_H Y).$$

Note that $\Omega^\alpha|_V$ lands in the right codomain because

$$R_Y \circ \pi \circ (\alpha_i|_V, \beta) = r_Y \circ \beta,$$

irrespective of $i \in \{1, 2\}$. We also note that the codomain of $\Omega^\alpha|_V$ is exactly the domain of the inverse $\Psi = (\Psi_1, \Psi_2)$ of the action map of the balanced tensor product $G \curvearrowright^{L_X} X \otimes_H Y \xrightarrow{R_Y} H_0$ (given explicitly in Proposition 5.20). In particular we then get a smooth map

$$\Psi_1 \circ \Omega^\alpha|_V : V \xrightarrow{\Omega^\alpha|_V} (X \otimes_H Y) \times_{G_0}^{R_Y, R_Y} (X \otimes_H Y) \xrightarrow{\Psi_1} G.$$

We now extend this map to the entire domain U_α , and show that it is independent on the choice of plot β . For that, pick two points $t, \bar{t} \in U_\alpha$, so that by subductiveness of the left moment map l_Y we can find two plots, $\beta : V \rightarrow Y$ and $\bar{\beta} : \bar{V} \rightarrow Y$, defined on open neighbourhoods of t and $\bar{t}$, respectively, such that $r_X \circ \alpha_i|_V = l_Y \circ \beta$ and $r_X \circ \alpha_i|_{\bar{V}} = l_Y \circ \bar{\beta}$. Following the above construction, we get two smooth maps:

$$\begin{aligned} \Omega^\alpha|_V : s &\longmapsto (\alpha_1|_V(s) \otimes \beta(s), \alpha_2|_V(s) \otimes \beta(s)), \\ \bar{\Omega}^\alpha|_{\bar{V}} : s &\longmapsto (\alpha_1|_{\bar{V}}(s) \otimes \bar{\beta}(s), \alpha_2|_{\bar{V}}(s) \otimes \bar{\beta}(s)). \end{aligned}$$

We now remark an important characterisation of Ψ , as a consequence of it being a diffeomorphism and inverse to the action map. Namely, when evaluated, $\Psi_1(x_1 \otimes y_1, x_2 \otimes y_2)$ is the *unique* arrow $g \in G$ satisfying the equation $gx_2 \otimes y_2 = x_1 \otimes y_1$. Therefore, $\Psi_1 \circ \Omega^\alpha|_V(s) \in G$ is the unique arrow such that

$$[\Psi_1 \circ \Omega^\alpha|_V(s)] \cdot (\alpha_2|_V(s) \otimes \beta(s)) = \alpha_1|_V(s) \otimes \beta(s).$$

By Lemma 5.27 all of the four actions of the original bibundles are free. Consequently, applying Lemma 5.28, since the second component in each term is just $\beta(s)$, this means that $\Psi_1 \circ \Omega^\alpha|_V(s)$ is the unique arrow in G such that

$$\Psi_1 \circ \Omega^\alpha|_V(s) \cdot \alpha_2|_V(s) = \alpha_1|_V(s),$$

where the tensor with $\beta(s)$ can be removed. But, for exactly the same reasons, if we take $s \in V \cap \bar{V}$, then $\Psi_1 \circ \bar{\Omega}^\alpha|_{\bar{V}}(s) \in G$ is *also* the unique arrow such that

$$\Psi_1 \circ \bar{\Omega}^\alpha|_{V \cap \bar{V}}(s) \cdot \alpha_2|_{V \cap \bar{V}}(s) = \alpha_1|_{V \cap \bar{V}}(s),$$

proving that

$$\Psi_1 \circ \Omega^\alpha|_{V \cap \bar{V}} = \Psi_1 \circ \bar{\Omega}^\alpha|_{V \cap \bar{V}}.$$

This shows that on the overlaps $V \cap \bar{V}$ the map $\Psi_1 \circ \Omega^\alpha|_{V \cap \bar{V}}$ does *not* depend on the plots β and $\bar{\beta}$. This allows us to extend $\Psi_1 \circ \Omega^\alpha|_V$, in a well-defined way, to the entire domain of U_α . We do this as follows. For every $t \in U_\alpha$ there exists a plot $\beta_t : V_t \rightarrow Y$, defined on an open neighbourhood $V_t \ni t$, such that $r_X \circ \alpha_i|_{V_t} = l_Y \circ \beta_t$. Clearly, this gives an open cover $(V_t)_{t \in U_\alpha}$ of U_α . For $t \in U_\alpha$ we then set $\Psi_1 \circ \Omega^\alpha(t) := \Psi_1 \circ \Omega^\alpha|_{V_t}(t)$. Hence we get a well-defined function $\Psi_1 \circ \Omega^\alpha : U_\alpha \rightarrow G$, which is smooth by the Axiom of Locality.

The main observation now is that, as the plot α is centred at (x_1, x_2) , we get that $\Psi_1 \circ \Omega^\alpha(0)$ is the unique arrow in G such that $\Psi_1 \circ \Omega^\alpha(0) \cdot x_2 = x_1$. This is exactly the property that characterises the division $\langle x_1, x_2 \rangle_G$!

Proposition 5.30. *A weakly invertible diffeological bibundle is pre-biprincipal.*

Proof. The bulk of the work has been done in Construction 5.29. Start with a diffeological bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ and a weak inverse $H \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} G$. We shall define a smooth division map $\langle \cdot, \cdot \rangle_G$ for the left G -action. For $(x_1, x_2) \in X \times_{H_0}^{r_X, r_X} X$, we know by the Axiom of Covering that the constant map $\text{const}_{(x_1, x_2)} : \mathbb{R} \rightarrow X \times_{H_0}^{r_X, r_X} X$ is a plot centred at (x_1, x_2) . We use the shorthand $\Psi_1 \circ \Omega^{(x_1, x_2)}$ to denote the map $\Psi_1 \circ \Omega^\alpha$ defined by the plot $\alpha = \text{const}_{(x_1, x_2)}$, and then write:

$$\langle x_1, x_2 \rangle_G := \Psi_1 \circ \Omega^{(x_1, x_2)}(0).$$

That just leaves us to show that this map is smooth. For that, take an arbitrary plot $\alpha : U_\alpha \rightarrow X \times_{H_0}^{r_X, r_X} X$ of the fibred product. We need to show that $\langle \cdot, \cdot \rangle_G \circ \alpha$ is a plot of G . For any $t \in U_\alpha$, we have that

$$\langle \alpha_1(t), \alpha_2(t) \rangle_G = \Psi_1 \circ \Omega^{\alpha(t)}(0)$$

is the unique arrow in G such that

$$\Psi_1 \circ \Omega^{\alpha(t)}(0) \cdot \text{const}_{\alpha(t)}^2(0) = \text{const}_{\alpha(t)}^1(0),$$

where const^i denotes the i th component of the constant plot. But then $\text{const}_{\alpha(t)}^i(0) = \alpha_i(t)$, and we already know that $\Psi_1 \circ \Omega^\alpha(t) \in G$ is the unique arrow that sends $\alpha_2(t)$ to $\alpha_1(t)$, so we have:

$$\Psi_1 \circ \Omega^{\alpha(t)}(0) = \Psi_1 \circ \Omega^\alpha(t), \quad \text{which means} \quad \langle \cdot, \cdot \rangle_G \circ \alpha = \Psi_1 \circ \Omega^\alpha.$$

But the right hand side $\Psi_1 \circ \Omega^\alpha : U_\alpha \rightarrow G$ is a plot of G as per Construction 5.29, proving that the map $\langle \cdot, \cdot \rangle_G$ is smooth. It is quite evident from its construction that it satisfies exactly the properties of a division map, and it is now easy to verify that

$$(\langle \cdot, \cdot \rangle_G, \text{pr}_2|_{X \times_{H_0} X}) : X \times_{H_0}^{r_X, r_X} X \longrightarrow G \times_{G_0}^{\text{src}, l_X} X$$

is a smooth inverse of the action map (see Section 4.2.1). The fact that it lands in the right codomain, i.e., $\text{src}(\langle x_1, x_2 \rangle_G) = l_X(x_2)$, follows from the properties of Ψ as the inverse of the action map of the balanced tensor product. Therefore $G \curvearrowright^{l_X} X \xrightarrow{r_X} H_0$ is a pre-principal bundle. An analogous argument will show that $G_0 \xleftarrow{l_Y} X \curvearrowright^{r_X} H$ is also pre-principal, and hence we have proved the claim. $\square$

We can now prove our main theorem:

Theorem 5.31. *A bibundle is weakly invertible in $\mathbf{DiffeolBiBund}$ if and only if it is biprincipal. That means: two diffeological groupoids are Morita equivalent if and only if they are equivalent in $\mathbf{DiffeolBiBund}$.*

Proof. One of the implications is just Proposition 5.24. The other now follows from a combination of Propositions 5.26 and 5.30. $\square$

This significantly generalises [Lan01b, Proposition 4.21], not only in that we have a generalisation to a diffeological setting, but also in that it considers a more general type of bibundle. It justifies the bicategory **DiffeolBiBund** as being the appropriate setting for Morita equivalence of diffeological groupoids. It also shows that the assumptions of left principality of the Lie groupoid bibundles appear to be more like technical necessities for getting a well defined bicategory of Lie groupoids and bibundles, rather than being meaningful assumptions on the underlying smooth structure of the bibundles. In Section 7.1 we discuss other aspects of diffeological Morita equivalence between Lie groupoids. A possible *category of fractions* approach to Morita equivalence of diffeological groupoids is discussed in [vdS20, Chapter V].

6. Some Morita Invariants

In theories of Morita equivalence, there are often interesting properties that are naturally Morita invariant. In this section we discuss some results that generalise several well known Morita invariants of Lie groupoids to the diffeological setting. These include: invariance of the orbit spaces (Definition 3.4), of being *fibrating* (Definition 6.2), and of the action categories (Definition 4.5). The proofs are taken from [vdS20, Chapter IV].

6.1 Invariance of orbit spaces

It is a well known result that if two Lie groupoids $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ are Morita equivalent (in the Lie groupoid sense), then there is a *homeomorphism* between their orbit spaces G_0/G and H_0/H , see e.g. [CM18, Lemma 1]. In fact, it turns out that the orbit spaces are even diffeologically diffeomorphic [Wat20, Theorem 3.8]⁸. The following theorem extends this result further from Lie groupoids to arbitrary diffeological groupoids. The construction of the underlying function is the same as for the Lie groupoid case, which is sketched in the proof of [CM18, Lemma 1], and which we describe below in detail.

⁸The author thanks the anonymous referee for bringing this result to his attention. We should also like to note that several other variants of this statement hold, namely in the settings of *differentiable*- and *subcartesian spaces*, as proved in [CM18].

Theorem 6.1. *If $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ are two Morita equivalent diffeological groupoids, then there is a diffeomorphism $G_0/G \cong H_0/H$ between their orbit spaces.*

Proof. Let $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ be the bibundle instantiating the Morita equivalence. Our first task will be to construct a function $\Phi : G_0/G \rightarrow H_0/H$ between the orbit spaces. The idea is to lift a point $a \in G_0$ of the base of the groupoid to its l_X -fibre, which by right principality is just an H -orbit in X , and then to project this orbit down to the other base H_0 along the right moment map r_X . The fact that the bundle is biprincipal ensures that this can be done in a consistent fashion.

We are dealing with *four* actions here, so we need to slightly modify our notation to avoid confusion. If $a \in G_0$ is an object in the groupoid G , we shall denote its orbit by $\text{Orb}_{G_0}(a)$, which, as usual, is just the set of all points $a' \in G_0$ such that there exists an arrow $g : a \rightarrow a'$ in G . Similarly, for $b \in H_0$ we write $\text{Orb}_{H_0}(b)$. On the other hand, we have two actions on X , for whose orbits we use the standard notations $\text{Orb}_G(x)$ and $\text{Orb}_H(x)$, where $x \in X$.

Now, start with a point $a \in G_0$, and consider its fibre $l_X^{-1}(a)$ in X . Since the bibundle is right subductive, the map l_X is surjective, so this fibre is non-empty and we can find a point $x_a \in l_X^{-1}(a)$. We claim that the expression $\text{Orb}_{H_0} \circ r_X(x_a)$ is independent on the choice of the point x_a in the fibre. For that, take another point $x'_a \in l_X^{-1}(a)$. This gives the equation $l_X(x_a) = l_X(x'_a)$, and since the bibundle is right pre-principal, we get a unique arrow $h \in H$ such that $x'_a = x_a h$. From the definition of a right groupoid action, this in turn gives the equations $r_X(x'_a) = \text{src}(h)$ and $r_X(x_a) = \text{trg}(h)$, which proves the claim. To summarise, whenever $x_a, x'_a \in l_X^{-1}(a)$ are two points in the same l_X -fibre, then we have:

$$\text{Orb}_{H_0} \circ r_X(x_a) = \text{Orb}_{H_0} \circ r_X(x'_a). \quad (1)$$

Next we want to show that neither is this expression dependent on the point $a \in G_0$, but rather on its orbit $\text{Orb}_{G_0}(a)$. For this, take another point $b \in \text{Orb}_{G_0}(a)$, so there exists some arrow $g : a \rightarrow b$ in G . Pick then $x \in l_X^{-1}(a)$ and $y \in l_X^{-1}(b)$. This means that $\text{src}(g) = l_X(x)$ and $\text{trg}(g) = l_X(y)$, which means that if we let g act on the point x we get a point $gx \in l_X^{-1}(b)$, in the same l_X -fibre as y . Then using equation (1) applied to gx and y , and

the G -invariance of the right moment map r_X , we immediately get:

$$\text{Orb}_{H_0} \circ r_X(x) = \text{Orb}_{H_0} \circ r_X(gx) = \text{Orb}_{H_0} \circ r_X(y).$$

Using this, we can now conclude that there is a well-defined function

$$\Phi : G_0/G \longrightarrow H_0/H; \quad \text{Orb}_{G_0}(a) \longmapsto \text{Orb}_{H_0} \circ r_X(x_a),$$

that is neither dependent on the point a in the orbit $\text{Orb}_{G_0}(a)$, nor on the choice of the point $x_a \in l_X^{-1}(a)$ in the fibre. Note that this function exists by virtue of right subductivity (and the Axiom of Choice), which ensures that the left moment map l_X is a surjection (and for each a there exists an x_a).

Either by replacing $G \curvearrowright^X X \curvearrowright^r H$ by its opposite bibundle, or by switching the words ‘left’ and ‘right’, the above argument analogously gives a function going the other way:

$$\Psi : H_0/H \longrightarrow G_0/G; \quad \text{Orb}_{H_0}(b) \longmapsto \text{Orb}_{G_0} \circ l_X(y_b),$$

where now $y_b \in r_X^{-1}(b)$ is some point in the fibre of the right moment map r_X . We claim that Φ and Ψ are mutual inverses. To see this, pick a point $a \in G_0$, a point $x_a \in l_X^{-1}(a)$, a point $y_{r_X(x_a)} \in r_X^{-1}(r_X(x_a))$. Then we can write

$$\Psi \circ \Phi (\text{Orb}_{G_0}(a)) = \Psi (\text{Orb}_{H_0}(r_X(x_a))) = \text{Orb}_{G_0} (l_X(y_{r_X(x_a)})).$$

We also have, by choice, the equation $r_X(x_a) = r_X(y_{r_X(x_a)})$, so by left pre-principality there exists an arrow $g \in G$ such that $gx_a = y_{r_X(x_a)}$. By definition of a left groupoid action, this then further gives

$$\text{src}(g) = l_X(x_a) = a \quad \text{and} \quad \text{trg}(g) = l_X(y_{r_X(x_a)}).$$

This proves that the right-hand side of the previous equation is equal to

$$\text{Orb}_{G_0} (l_X(y_{r_X(x_a)})) = \text{Orb}_{G_0}(a),$$

which gives $\Psi \circ \Phi = \text{id}_{G_0/G}$. Through a similar argument, using right pre-principality, we obtain that $\Phi \circ \Psi = \text{id}_{H_0/H}$.

To finish the proof, it suffices to prove that both Φ and Ψ are smooth. Again, due to the symmetry of the situation, we shall only prove that Φ is

smooth. The proof for Ψ will follow analogously. Since Orb_{G_0} is a subduction, to prove that Φ is smooth it suffices by Lemma 2.20(3) to prove that $\Phi \circ \text{Orb}_{G_0}$ is smooth. Since the left moment map l_X is a surjection, using the Axiom of Choice we pick a section $\sigma : G_0 \rightarrow X$, which replaces our earlier notation of $\sigma(a) =: x_a$. From the way Φ is defined, we see that we get a commutative diagram:

$$\begin{array}{ccccc} G_0 & \xrightarrow{\sigma} & X & \xrightarrow{r_X} & H_0 \\ \text{Orb}_{G_0} \downarrow & & & & \downarrow \text{Orb}_{H_0} \\ G_0/G & \xrightarrow{\Phi} & & & H_0/H. \end{array}$$

We are therefore to show that $\text{Orb}_{H_0} \circ r_X \circ \sigma$ is smooth. For this, pick a plot $\alpha : U_\alpha \rightarrow G_0$ of the base space. By right subductivity, the left moment map l_X is a subduction, so locally $\alpha|_V = l_X \circ \beta$, where β is some plot of X . Now, note that, for all $t \in V$, both the points $\beta(t)$ and $\sigma \circ l_X \circ \beta(t)$ are elements of the fibre $l_X^{-1}(l_X \circ \beta(t))$. Therefore, by equation (1) we get:

$$\text{Orb}_{H_0} \circ r_X \circ \sigma \circ \alpha|_V = \text{Orb}_{H_0} \circ r_X \circ \sigma \circ l_X \circ \beta = \text{Orb}_{H_0} \circ r_X \circ \beta.$$

The right-hand side of this equation is clearly smooth (and no longer dependent on the choice of section σ). By the Axiom of Locality for G_0 , it follows that $\text{Orb}_{H_0} \circ r_X \circ \sigma \circ \alpha$ is globally smooth, and since the plot α was arbitrary, this proves that $\Phi \circ \text{Orb}_{G_0}$ is smooth. Hence, Φ is smooth. After an analogous argument that shows Ψ is smooth, the desired diffeomorphism between the orbit spaces follows. $\square$

Note that in the proof of [Wat20, Theorem 3.8], instead of a global (not necessarily smooth) section $\sigma : G_0 \rightarrow X$ of the left moment map $l_X : X \rightarrow G_0$, they use the fact that l_X is a surjective submersion to find a *local smooth* section. Our proof shows that it is not necessary for σ to be smooth, highlighting another difference between the rôle of surjective submersions and subductions⁹.

6.2 Invariance of fibration

The theory of diffeological (principal) fibre bundles is shown in [IZ13a, Chapter 8] to be fully captured by the following notion:

⁹We thank the anonymous referee for pointing out this difference between the proofs.

Definition 6.2. A diffeological groupoid $G \rightrightarrows G_0$ is called *fibrating* (or a *fibration groupoid*) if the characteristic map $(\text{trg}, \text{src}) : G \rightarrow G_0 \times G_0$ is a subduction.

This leads to a theory of diffeological fibre bundles that is able to treat the standard smooth locally trivial (principal) fibre bundles of smooth manifolds, but also bundles that are not (and could not meaningfully be) locally trivial. It is then natural to ask if this property of diffeological groupoids is invariant under Morita equivalence. The following theorem proves that this is the case:

Theorem 6.3. Let $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ be two Morita equivalent diffeological groupoids. Then $G \rightrightarrows G_0$ is fibrating if and only if $H \rightrightarrows H_0$ is fibrating.

Proof. Because Morita equivalence is an equivalence relation, it suffices to prove that if $G \rightrightarrows G_0$ is fibrating, then so is $H \rightrightarrows H_0$. Denoting the characteristic maps of these groupoids by $\chi_G = (\text{trg}_G, \text{src}_G)$ and $\chi_H = (\text{trg}_H, \text{src}_H)$, assume that G is fibrating, so that χ_G is a subduction. Our goal is to show χ_H is also a subduction.

To begin with, take an arbitrary plot $\alpha = (\alpha_1, \alpha_2) : U_\alpha \rightarrow H_0 \times H_0$, and fix an element $t \in U_\alpha$. We thus need to find a plot $\Phi : W \rightarrow H$, defined on an open neighbourhood $t \in W \subseteq U_\alpha$, such that $\alpha|_W = \chi_H \circ \Phi$. Morita equivalence yields a biprincipal bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$. To construct the plot Φ , we use almost all of the structure of this bibundle.

The right moment map $r_X : X \rightarrow H_0$ is a subduction, so for each of the components α_i of α we get a plot $\beta_i : U_i \rightarrow X$, defined on an open neighbourhood $t \in U_i \subseteq U_\alpha$, such that $\alpha_i|_{U_i} = r_X \circ \beta_i$. Define $U := U_1 \cap U_2$, which is another open neighbourhood of $t \in U_\alpha$, and introduce the notation

$$\beta := (\beta_1|_U, \beta_2|_U) : U \longrightarrow X \times X.$$

Composing with the left moment map $l_X : X \rightarrow G_0$, we get $(l_X \times l_X) \circ \beta : U \rightarrow G_0 \times G_0$. It is here that we use that $G \rightrightarrows G_0$ is fibrating. Because of that, we can find an open neighbourhood $t \in V \subseteq U \subseteq U_\alpha$ and a plot $\Omega : V \rightarrow G$ such that

$$\chi_G \circ \Omega = (l_X \times l_X) \circ \beta|_V. \quad (2)$$

This means that $\text{trg}_G \circ \Omega = l_X \circ \beta_1|_V$ and $\text{src}_G \circ \Omega = l_X \circ \beta_2|_V$. Let $\varphi_G : X \otimes_H \overline{X} \rightarrow G$ be the biequivariant diffeomorphism from Proposition 5.24. Using the plot Ω we just obtained, we get another plot $\varphi_G^{-1} \circ \Omega : V \rightarrow X \otimes_H \overline{X}$. Now, since the canonical projection $\pi_H : X \times_{H_0}^{r_X, r_X} \overline{X} \rightarrow X \otimes_H \overline{X}$ of the diagonal H -action is a subduction, we can find an open neighbourhood $t \in W \subseteq V$ and a plot $\omega : W \rightarrow X \times_{H_0}^{r_X, r_X} \overline{X}$ such that

$$\pi_H \circ \omega = \varphi_G^{-1} \circ \Omega|_W. \quad (3)$$

Note that the plot ω decomposes into its components $\omega_1, \omega_2 : W \rightarrow X$, which satisfy $r_X \circ \omega_1 = r_X \circ \omega_2$. Using the biequivariance of φ_G and the defining relation $L_X \circ \pi_H = l_X \circ \text{pr}_1|_{X \times_{H_0} \overline{X}}$ we find:

$$\begin{aligned} l_X \circ \beta_1|_W &= \text{trg}_G \circ \Omega|_W \\ &= L_X \circ \varphi_G^{-1} \circ \Omega|_W \\ &= L_X \circ \pi_H \circ \omega \\ &= l_X \circ \text{pr}_1|_{X \times_{H_0} \overline{X}} \circ \omega \\ &= l_X \circ \omega_1, \end{aligned}$$

where the first equality follows from the equation (2), and the third one from (3). Similarly, we find $l_X \circ \beta_2|_W = l_X \circ \omega_2$. These two equalities give two well-defined plots, one for each $i \in \{1, 2\}$, given by

$$\beta_i|_W \otimes \omega_i := \pi_G \circ (\beta_i|_W, \omega_i) : W \xrightarrow{(\beta_i|_W, \omega_i)} \overline{X} \times_{G_0}^{l_X, l_X} X \xrightarrow{\pi_G} \overline{X} \otimes_G X,$$

where $\pi_G : \overline{X} \times_{G_0}^{l_X, l_X} X \rightarrow \overline{X} \otimes_G X$ is the canonical projection of the diagonal G -action. We can now apply the biequivariant diffeomorphism $\varphi_H : \overline{X} \otimes_G X \Rightarrow H$ from Proposition 5.24 to get two plots in H . It is from these two plots that we will create Φ . Here it is absolutely essential that we have constructed the plot ω such that $r_X \circ \omega_1 = r_X \circ \omega_2$, because that means that the sources of these two plots in H will be equal, and hence they can be composed if we first invert one of them component-wise. To see this, use the biequivariance of φ_H to calculate

$$\begin{aligned} \text{src}_H \circ \varphi_H \circ (\beta_i|_W \otimes \omega_i) &= R_X \circ (\beta_i|_W \otimes \omega_i) \\ &= r_X \circ \text{pr}_2|_{\overline{X} \times_{G_0} X} \circ (\beta_i|_W, \omega_i) \\ &= r_X \circ \omega_i, \end{aligned}$$

and similarly:

$$\begin{aligned}
 \mathrm{trg}_H \circ \varphi_H \circ (\beta_i|_W \otimes \omega_i) &= L_{\overline{X}} \circ (\beta_i|_W \otimes \omega_i) \\
 &= r_X \circ \mathrm{pr}_1|_{\overline{X} \times_{G_0} X} \circ (\beta_i|_W, \omega_i) \\
 &= r_X \circ \beta_i|_W \\
 &= \alpha_i|_W.
 \end{aligned}$$

Of course, if we switch $\beta_i|_W \otimes \omega_i$ to $\omega_i \otimes \beta_i|_W$, which is defined in the obvious way, then the right-hand sides of the above two equations will switch. So, for every $s \in W$, the expression $\varphi_H(\omega_2(s) \otimes \beta_2(s))$ is an arrow in H from $r_X \circ \beta_2(s) = \alpha_2(s)$ to $r_X \circ \omega_2(s)$, and $\varphi_H(\beta_1(s) \otimes \omega_1(s))$ is an arrow from $r_X \circ \omega_1(s) = r_X \circ \omega_2(s)$ to $r_X \circ \beta_1(s) = \alpha_1(s)$, which can hence be composed to give an arrow from $\alpha_2(s)$ to $\alpha_1(s)$. This is exactly the kind of arrow we want. Therefore, for every $s \in W$, we get a commutative triangle in the groupoid H , which defines for us the plot $\Phi : W \rightarrow H$:

$$\begin{array}{ccc}
 \alpha_2(s) & \xrightarrow{\quad \Phi(s) \quad} & \alpha_1(s) \\
 \searrow \varphi_H(\omega_2(s) \otimes \beta_2(s)) & & \nearrow \varphi_H(\beta_1(s) \otimes \omega_1(s)) \\
 & r_X \circ \omega_1(s) &
 \end{array}$$

The map Φ is clearly smooth, because inversion and multiplication in H are smooth. Hence we have defined the plot Φ , and by the above diagram it is clear that it satisfies

$$\chi_H \circ \Phi = (\mathrm{trg}_H \circ \Phi, \mathrm{src}_H \circ \Phi) = \alpha|_W.$$

Thus we may at last conclude that χ_H is a subduction, and hence that $H \rightrightarrows H_0$ is also fibrating. $\square$

6.3 Invariance of representations

In the Morita theory of rings, it holds that two rings are Morita equivalent if and only if their categories of modules are equivalent. For groupoids, even discrete ones, this is no longer an “if and only if” proposition, but merely an “only if”. Nevertheless, it is known that the result transfers to Lie groupoids as well [Lan01a, Theorem 6.6], and here we shall prove that it transfers also to diffeology.

Theorem 6.4. *Suppose that $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ are Morita equivalent diffeological groupoids. Then the action categories $\mathbf{Act}(G \rightrightarrows G_0)$ and $\mathbf{Act}(H \rightrightarrows H_0)$ are categorically equivalent.*

Proof. If $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ are Morita equivalent, there exists a bi-principal bibundle $G \curvearrowright^{L_X} X \curvearrowright^{R_X} H$. Recall from Definition 4.5 the notion of action categories and from Definition 5.9 that of induced action functors. We claim that

$$\begin{aligned} X \otimes_H - : \mathbf{Act}(H \rightrightarrows H_0) &\longrightarrow \mathbf{Act}(G \rightrightarrows G_0), \\ \overline{X} \otimes_G - : \mathbf{Act}(G \rightrightarrows G_0) &\longrightarrow \mathbf{Act}(H \rightrightarrows H_0) \end{aligned}$$

are mutually inverse functors up to natural isomorphism. To see this, take a left H action $H \curvearrowright^{L_Y} Y$. Then

$$\begin{aligned} (\overline{X} \otimes_G -) \circ (X \otimes_H -) [H \curvearrowright^{L_Y} Y] &= (\overline{X} \otimes_G -) [G \curvearrowright^{L_X} X \otimes_H Y] \\ &= H \curvearrowright^{L_{\overline{X}}} (\overline{X} \otimes_G (X \otimes_H Y)). \end{aligned}$$

Therefore, we need to construct a natural biequivariant diffeomorphism

$$\mu_Y : \overline{X} \otimes_G (X \otimes_H Y) \longrightarrow Y.$$

For this, we collect the biequivariant diffeomorphisms from Propositions 5.12, 5.13 and 5.24. Let us denote them by

$$\begin{aligned} A_Y : \overline{X} \otimes_G (X \otimes_H Y) &\longrightarrow (\overline{X} \otimes_G X) \otimes_H Y, \\ \varphi_H : \overline{X} \otimes_G X &\longrightarrow H, \\ M_Y : H \otimes_H Y &\longrightarrow Y, \end{aligned}$$

describing the association up to isomorphism, the division map of the bibundle, and the left action $H \curvearrowright Y$, respectively. We then define

$$\mu_Y := M_Y \circ (\varphi_H \otimes \text{id}_Y) \circ A_Y.$$

Note that $(\varphi_H \otimes \text{id}_Y)$ is still a biequivariant diffeomorphism. The naturality square of the natural transformation $\mu : (\overline{X} \otimes_G -) \circ (X \otimes_H -) \Rightarrow \text{id}_{\mathbf{Act}(H)}$ then becomes:

$$\begin{array}{ccc} \overline{X} \otimes_G (X \otimes_H Y) & \xrightarrow{\mu_Y} & Y \\ \text{id}_{\overline{X}} \otimes (\text{id}_X \circ \varphi) \downarrow & & \downarrow \varphi \\ \overline{X} \otimes_G (X \otimes_H Z) & \xrightarrow{\mu_Z} & Z, \end{array}$$

where $\varphi : Y \rightarrow Z$ is an H -equivariant smooth map. It follows from the structure of these maps that the naturality square commutes. The top right corner of the diagram becomes:

$$\begin{aligned}
 \varphi \circ \mu_Y (x_1 \otimes (x_2 \otimes y)) &= \varphi \circ M_Y \circ (\varphi_H \otimes \text{id}_Y) \circ A_Y (x_1 \otimes (x_2 \otimes y)) \\
 &= \varphi \circ M_Y \circ (\varphi_H \otimes \text{id}_Y) ((x_1 \otimes x_2) \otimes y) \\
 &= \varphi \circ M_Y (\varphi_H(x_1 \otimes x_2) \otimes y) \\
 &= \varphi (\varphi_H(x_1 \otimes x_2)y) \\
 &= \varphi_H(x_1 \otimes x_2)\varphi(y),
 \end{aligned}$$

where the very last step follows from H -equivariance of φ . Following a similar calculation, the bottom left corner evaluates as

$$\begin{aligned}
 \mu_Z \circ (\text{id}_{\overline{X}} \otimes (\text{id}_X \otimes \varphi)) &= M_Z \circ (\varphi_H \otimes \text{id}_Z) \circ A_Z \circ (\text{id}_{\overline{X}} \otimes (\text{id}_X \otimes \varphi)) \\
 &= M_Z \circ (\varphi_H \otimes \text{id}_Z) \circ ((\text{id}_{\overline{X}} \otimes \text{id}_X) \otimes \varphi) \\
 &= M_Z \circ (\varphi_H \otimes \varphi),
 \end{aligned}$$

which, when evaluated, gives exactly the same as the above expression for the top right corner. This proves that μ is natural, and since every of its components is an H -equivariant diffeomorphism, it follows that μ is a natural isomorphism. The fact that the composition $(X \otimes_H -) \circ (\overline{X} \otimes_G -)$ is naturally isomorphic to $\text{id}_{\text{Act}(G)}$ follows from an analogous argument. Hence the categories $\text{Act}(G \rightrightarrows G_0)$ and $\text{Act}(H \rightrightarrows H_0)$ are equivalent, as was to be shown. $\square$

7. Discussion and Suggestions for Future Research

7.1 Diffeological bibundles between Lie groupoids

As we saw in Example 5.4, if two Lie groupoids are *Lie* Morita equivalent (i.e. Morita equivalent in the Lie groupoid sense [CM18, Definition 2.15]), then they are also *diffeologically* Morita equivalent. This is simply due to the fact that surjective submersions between smooth manifolds are in particular also subductions, and hence a Lie principal groupoid bundle is also diffeologically principal. But, what if $G \rightrightarrows G_0$ and $H \rightrightarrows H_0$ are two *Lie* groupoids, such that there exists a *diffeological* biprincipal bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$

between them. What does that say about the *Lie* Morita equivalence of G and H ? This still remains an open question (Question 7.6). In this section we discuss some related results, which also pertain to our choice of subductions over *local* subductions for the development of the general theory. A slightly more detailed discussion is in [vdS20, Section 4.4.3]. In light of Proposition 2.25, the source and target maps of a Lie groupoid are local subductions (cf. Proposition 3.2), and we can therefore introduce the following class of diffeological groupoids:

Definition 7.1. *We say a diffeological groupoid $G \rightrightarrows G_0$ is locally subductive if its source and target maps are local subductions¹⁰. Clearly, every Lie groupoid is a locally subductive diffeological groupoid.*

Looking at the structure of the proofs in Sections 4 and 5, it appears as if they can be generalised to a setting where we replace all subductions by local subductions. In doing so, we would get a theory of locally subductive groupoids, locally subductive groupoid bundles, and the corresponding notions for bibundles and Morita equivalence, which, as it appears, would follow the same story as we have so far presented. An upside to that framework would be that it directly returns the original theory of Morita equivalence for Lie groupoids, once we restrict our diffeological spaces to smooth manifolds. In this section we shall prove that, even in the slightly more general setting of Section 5, the diffeological bibundle theory reduces to the Lie groupoid theory in the correct way. We do this by proving that the moment maps of a biprincipal bibundle between locally subductive groupoids have to be local subductions as well (Lemma 7.3). In hindsight, this provides more justification for our choice of starting with subductions instead of local subductions. One consequence of this choice is that it allows for groupoid bundles that are truly *pseudo*-bundles, in the sense of [Per16]. The notion of pseudo-bundles seems to be the correct notion in the setting of diffeology to generalise all bundle constructions on manifolds, at least if we want to treat (internal) tangent bundles as such (see [CW16]). There exists diffeological spaces whose internal tangent bundle is not a local subduction [CW16, Example 3.17]. If we had defined principality of a groupoid bundle to include

¹⁰It would be tempting to call such groupoids “*diffeological Lie groupoids*,” but this would conflict with earlier established terminology of so-called *diffeological Lie groups* in [IZ13a, Article 7.1] and [Les03; Mag18].

local subductiveness, these examples would not be treatable by our theory of Morita equivalence.

Lemma 7.2. *Let $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ be a diffeological bibundle, where $H \rightrightarrows H_0$ is a locally subductive groupoid. Then the canonical quotient projection map $\pi_H : X \times_{H_0}^{r_X, r_X} \overline{X} \rightarrow X \otimes_H \overline{X}$ is a local subduction.*

Proof. Let $\alpha : (U_\alpha, 0) \rightarrow (X \otimes_H \overline{X}, x_1 \otimes x_2)$ be a pointed plot of the balanced tensor product. Since π_H is already a subduction, we can find a plot $\beta : V \rightarrow X \times_{H_0} \overline{X}$, defined on an open neighbourhood $0 \in V \subseteq U_\alpha$ of the origin, such that $\alpha|_V = \pi_H \circ \beta$. This plot decomposes into two plots $\beta_1, \beta_2 \in \mathcal{D}_X$ on X , satisfying $r_X \circ \beta_1 = r_X \circ \beta_2$. We use the notation $\alpha|_V = \beta_1 \otimes \beta_2$. In particular, we get an equality $x_1 \otimes x_2 = \beta_1(0) \otimes \beta_2(0)$ inside the balanced tensor product, which means that we can find an arrow $h \in H$ such that $\beta_i(0) = x_i h$. The target must be $\text{trg}(h) = r_X(x_1) = r_X(x_2)$. This arrow allows us to write a pointed plot $r_X \circ \beta_i : (V, 0) \rightarrow (H_0, \text{trg}(h^{-1}))$, so that now we can use that $H \rightrightarrows H_0$ is locally subductive. Since the target map of H is a local subduction, we can find a pointed plot $\Omega : (W, 0) \rightarrow (H, h^{-1})$ such that $r_X \circ \beta_i|_W = \text{trg}_H \circ \Omega$. This relation means that, for every $t \in W$, we have a well-defined action $\beta_i(t) \cdot \Omega(t) \in X$. Hence we get a pointed plot

$$\Psi : (W, 0) \longrightarrow (X \times_{H_0}^{r_X, r_X} \overline{X}, (x_1, x_2)); \quad t \longmapsto (\beta_1(t)\Omega(t), \beta_2(t)\Omega(t)).$$

It then follows by the definition of the balanced tensor product that

$$\pi_H \circ \Psi(t) = \beta_1|_W(t)\Omega(t) \otimes \beta_2|_W(t)\Omega(t) = \beta_1|_W(t) \otimes \beta_2|_W(t) = \alpha|_W(t),$$

proving that π_H is a local subduction. $\square$

Lemma 7.3. *If $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ is a biprincipal bibundle between locally subductive groupoids, then the moment maps l_X and r_X are local subductions as well.*

Proof. If the bibundle $G \curvearrowright^{l_X} X \curvearrowright^{r_X} H$ is biprincipal, we get two biequivariant diffeomorphisms $\varphi_G : X \otimes_H \overline{X} \rightarrow G$ and $\varphi_H : \overline{X} \otimes_G X \rightarrow H$ (Proposition 5.24). It follows that the local subductivity of the source and target maps of G and H transfer to the four moment maps of the balanced tensor products. For example, the left moment map $L_X : X \otimes_H \overline{X} \rightarrow G_0$ can

be written as $L_X = \text{trg}_G \circ \varphi_G$, where the right hand side is clearly a local subduction. We know as well that L_X fits into a commutative square with the original moment map l_X :

$$\begin{array}{ccc} X \times_{H_0}^{r_X, r_X} \overline{X} & \xrightarrow{\pi_H} & X \otimes_H \overline{X} \\ \text{pr}_1|_{X \times_{H_0} \overline{X}} \downarrow & & \downarrow L_X \\ X & \xrightarrow{l_X} & G_0. \end{array}$$

Since local subductions compose, and since by Lemma 7.2 the projection π_H is a local subduction, we find that the upper right corner $L_X \circ \pi_H$ must be a local subduction. Hence the composition $l_X \circ \text{pr}_1|_{X \times_{H_0} \overline{X}}$ is a local subduction, which by an argument that is analogous to the proof of Lemma 2.20(2) gives the local subductiveness of l_X . That the right moment map r_X is a local subduction follows from a similar argument. $\square$

The lemma suggests that, if we refine our notion of principality to something we might call *pure-principality*, by passing from subductions to local subductions, then biprincipality between locally subductive groupoids means the same thing as this new notion of pure-principality. Let us make this precise.

Definition 7.4. *Two diffeological groupoids are called purely Morita equivalent if there exists a biprincipal bibundle between them, such that the two underlying moment maps are local subductions.*

Clearly, pure Morita equivalence implies ordinary Morita equivalence in the sense of Definition 5.3, since local subductions are, in particular, subductions. The question is if the converse implication holds as well. We have a partial answer, since Lemma 7.3 can now be restated as follows:

Proposition 7.5. *Two locally subductive groupoids are Morita equivalent if and only if they are purely Morita equivalent.*

Especially in light of the existence of subductions that are not local subductions (see e.g. [IZ13a, Exercise 61, p.60]), and the fact that the proof of Lemma 7.3 relies so heavily on the assumption that the groupoids are locally subductive, it seems that the ordinary diffeological Morita equivalence

of Definition 5.3 is not equivalent to pure-Morita equivalence in general. We do not, however, know of an explicit counter-example. This discussion leaves us an open question:

Question 7.6. *Does diffeological Morita equivalence reduce to Lie Morita equivalence on Lie groupoids? That is to ask, if two Lie groupoids are diffeologically Morita equivalent, are they also Lie Morita equivalent?*

If two Lie groupoids G and H are diffeologically Morita equivalent, then there exists a diffeological biprincipal bibundle $G \curvearrowright^L X \curvearrowright^R H$, where X is a diffeological space. A positive answer to Question 7.6 could consist of a proof that X is in fact a smooth manifold. Since G and H are both manifolds, it follows that $X \otimes_H \overline{X}$ and $\overline{X} \otimes_G X$ are also manifolds. We do not know if this is sufficient to imply that X itself has to be a manifold. One suggestion is to use [IZ13a, Article 4.6], which gives a characterisation for when a quotient of a diffeological space by an equivalence relation is a smooth manifold. Since the balanced tensor products are quotients of diffeological spaces, one may try to use this result to obtain a special family of plots for their underlying fibred products. This could potentially be used to define an atlas on X .

7.2 Directions for future research

We list here some possible directions for future research. These are also proposed at the end of [vdS20, Section 1.2.3].

- Finding an answer to the open Question 7.6 about *diffeological* Morita equivalence between *Lie* groupoids.
- The construction of a theory of bibundles for a more general framework of generalised smooth spaces. One possibility is to look at the *generalised spaces* of [BH11, Definition 4.11] (subsuming diffeology), or even to look at arbitrary classes of sheaves. What is the relation between our theory of Morita equivalence and the discussion in [MZ15]? A theory of principal bibundles seems to exist in a general setting for groupoids in ∞ -toposes: [nL18].
- What is the precise relation between differentiable stacks and diffeological groupoids (cf. [WW19])? Using our notion of Morita equi-

valence, what types of objects are “*diffeological stacks*” (i.e., Morita equivalence classes of diffeological groupoids)?

- Can the *Hausdorff Morita equivalence* for holonomy groupoids of singular foliations introduced in [GZ19] be understood as a Morita equivalence between diffeological groupoids?
- Can the bridge between diffeology and noncommutative geometry that is being built in [Ber16; IZL18; ASZ19; IZP20] be strengthened by our theory of Morita equivalence? Morita equivalence of Lie groupoids is already an important concept in relation to noncommutative geometry, especially for the theory of groupoid C^* -algebras. Can this link be extended to the diffeological setting, possibly through a theory of groupoid C^* -algebras for (a large class of) diffeological groupoids? If such a theory exists, what is the relation between Morita equivalence of diffeological groupoids and the Morita equivalence of their groupoid C^* -algebras? Is Morita equivalence preserved just like in the Lie case?

References

- [ASZ19] I. Androulidakis, G. Skandalis and M. Zambon. *Diffeological groupoids and their C^* -algebras*. Lecture slides, <https://indico.math.cnrs.fr/event/3571/contributions/1977/attachments/2341/2854/Androulidakis.pdf>. 2019.
- [AZ] I. Androulidakis and M. Zambon. *Integration of Singular Subalgebroids*. In preparation.
- [BH11] J. Baez and A. Hoffnung. “Convenient categories of smooth spaces”. In: *Transactions of the American Mathematical Society* 363.11 (2011), pp. 5789–5825.
- [BIKW17] A. Batubenge, P. Iglesias-Zemmour, Y. Karshon and J. Watts. *Diffeological, Frölicher, and Differential Spaces*. 2017. arXiv: 1712.04576 [math.DG].

- [Ber16] P. Bertozzini. *Spectral Geometries for Some Diffeologies*. Lecture slides, http://bcc.impan.pl/16Index-SIII/uploads/Bertozzini_slides.pdf. 2016.
- [Blo08] C. Blohmann. “Stacky Lie groups”. In: *Int. Math. Res. Not.* (2008), Art. ID rnn 082, 51.
- [BFW13] C. Blohmann, M. C. B. Fernandes and A. Weinstein. “Groupoid symmetry and constraints in general relativity”. In: *Communications in Contemporary Mathematics* 15.01 (2013), pp. 125–161.
- [Bos07] R. D. Bos. “Groupoids in Geometric Quantization”. PhD thesis. Radboud University Nijmegen, 2007.
- [CW16] J. Christensen and E. Wu. “Tangent spaces and tangent bundles for diffeological spaces”. In: *Cah. Topol. Géom. Différ. Catég. LVII* (2016).
- [CLW16] B. Collier, E. Lerman and S. Wolbert. “Parallel transport on principal bundles over stacks”. In: *Journal of Geometry and Physics* 107 (2016), pp. 187–213.
- [CM18] M. Crainic and J. N. Mestre. “Orbispace as differentiable stratified spaces”. In: *Letters in mathematical physics* 108.3 (2018), pp. 805–859.
- [dHo12] M. L. del Hoyo. *Lie Groupoids and Differentiable Stacks*. Dec. 2012. arXiv: 1212.6714 [math.DG].
- [Don84] P. Donato. “Revêtements et groupe fondamental des espaces différentiels homogènes”. PhD thesis. Université de Provence, Marseille, 1984.
- [DI83] P. Donato and P. Iglesias. “Exemple de groupes différentiels: flots irrationnels sur le tore”. In: *CPT-83/P-1524* (1983). <http://math.huji.ac.il/~piz/documents/EDGDFISLT.pdf>.
- [GZ19] A. Garmendia and M. Zambon. “Hausdorff Morita equivalence of singular foliations”. In: *Annals of Global Analysis and Geometry* 55.1 (2019), pp. 99–132.

- [Gl19] J. Głowacki. “Groupoid Symmetry and Constraints Bracket of General Relativity Revisited”. Master thesis, <https://www.math.ru.nl/~landsman/Jan2019.pdf>. 2019.
- [HS87] M. Hilsum and G. Skandalis. “Morphismes K -orientés d’espaces de feuilles et fonctorialité en théorie de Kasparov (d’après une conjecture d’A. Connes)”. In: *Annales scientifiques de l’École normale supérieure*. Vol. 20. 3. 1987, pp. 325–390.
- [Igl85] P. Iglesias. “Fibrations difféologiques et homotopie”. PhD thesis. Université de Provence, Marseille, 1985.
- [IZ13a] P. Iglesias-Zemmour. *Diffeology*. Vol. 185. American Mathematical Soc., 2013.
- [IZ13b] P. Iglesias-Zemmour. *The Beginning of Diffeological Spaces*. Blog post, <http://math.huji.ac.il/~piz/documents/DBlog-Rmk-TBODS.pdf>. 2013.
- [IZ17] P. Iglesias-Zemmour. *An Introduction to Diffeology*. <http://math.huji.ac.il/~piz/documents/AITD.pdf>. 2017.
- [IZL18] P. Iglesias-Zemmour and J.-P. Laffineur. “Noncommutative geometry and diffeology: The case of orbifolds”. In: *Journal of Noncommutative Geometry* 12.4 (2018), pp. 1551–1572.
- [IZP20] P. Iglesias-Zemmour and E. Prato. *Quasifolds, Diffeology and Noncommutative Geometry*. May 2020. arXiv: 2005.09283 [math.DG].
- [Lac10] S. Lack. “A 2-categories companion”. In: *Towards higher categories*. Springer, 2010, pp. 105–191.
- [Lan01a] N. P. Landsman. “Bicategories of operator algebras and Poisson manifolds”. In: *Mathematical physics in mathematics and physics*. Vol. 30. Fields Inst. Commun. Amer. Math. Soc., 2001, pp. 271–286.
- [Lan01b] N. P. Landsman. “Quantized reduction as a tensor product”. In: *Quantization of singular symplectic quotients*. Springer, 2001, pp. 137–180.

- [Les03] J. Leslie. “On a diffeological group realization of certain generalized symmetrizable Kac-Moody Lie algebras”. In: *Journal of Lie Theory* 13.2 (2003), pp. 427–442.
- [Mac71] S. Mac Lane. *Categories for the working mathematician*. Second Edition (1998). Vol. 5. Graduate Texts in Mathematics. Springer-Verlag, 1971.
- [Mac20] L. E. MacDonald. *The holonomy groupoids of singularly foliated bundles*. June 2020. arXiv: 2006.14271 [math.DG].
- [Mac05] K. C. H. Mackenzie. *General theory of Lie groupoids and Lie algebroids*. Vol. 213. Cambridge University Press, 2005.
- [Mag18] J.-P. Magnot. “The group of diffeomorphisms of a non-compact manifold is not regular”. In: *Demonstratio Mathematica* 51.1 (2018), pp. 8–16.
- [MZ15] R. Meyer and C. Zhu. “Groupoids in Categories with Pretopology”. In: *Theory & Applications of Categories* 30.55 (2015).
- [MM05] I. Moerdijk and J. Mrčun. “Lie groupoids, sheaves and cohomology”. In: *Poisson geometry, deformation quantisation and group representations*. Vol. 323. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 2005, pp. 145–272.
- [MRW87] P. S. Muhly, J. N. Renault and D. P. Williams. “Equivalence and isomorphism for groupoid C^* -algebras”. In: *Journal of Operator Theory* (1987), pp. 3–22.
- [nL18] nLab. *Bibundle*. <https://ncatlab.org/nlab/show/bibundle>, (version: 2018-07-3). 2018.
- [Per16] E. Pervova. “Diffeological vector pseudo-bundles”. In: *Topology and its Applications* 202 (2016), pp. 269–300.
- [Rie81] M. A. Rieffel. “ C^* -algebras associated with irrational rotations”. In: *Pacific Journal of Mathematics* 93.2 (1981), pp. 415–429.
- [RV18] D. M. Roberts and R. F. Vozzo. “Smooth loop stacks of differentiable stacks and gerbes”. In: *Cah. Topol. Géom. Différ. Catég.* LIX-2 (2018), pp. 95–141.

- [Ser65] J.-P. Serre. “Lie groups and Lie algebras”. In: *Lecture Notes in Math* 1500 (1965).
- [Sou80] J.-M. Souriau. “Groupes différentiels”. In: *Differential geometrical methods in mathematical physics*. Springer, 1980, pp. 91–128.
- [Sou84] J.-M. Souriau. “Groupes différentiels et physique mathématique”. In: *Group Theoretical Methods in Physics*. Springer, 1984, pp. 511–513.
- [Sta11] A. Stacey. “Comparative smootheology”. In: *Theory and Applications of Categories* 25.4 (2011), pp. 64–117.
- [vdS20] N. van der Schaaf. “Diffeology, Groupoids & Morita Equivalence”. Master Thesis, <https://www.math.ru.nl/~landsman/NestaM.pdf>. 2020.
- [Wat20] J. Watts. *The Orbit Space and Basic Forms of a Proper Lie Groupoid*. May 2020. arXiv: 1309.3001v4 [math.DG].
- [WW19] J. Watts and S. Wolbert. *Diffeological Coarse Moduli Spaces of Stacks over Manifolds*. 2019. arXiv: 1406.1392 [math.DG].

Nesta van der Schaaf
School of Informatics
University of Edinburgh
10 Crichton St, Newington
Edinburgh EH8 9AB (Scotland)
n.schaaf@ed.ac.uk